



PROCEEDINGS

of the


California Academy of Sciences

(Series 4)



December 31, 2015 * Volume 62 * Numbers 18–21

Institute for Biodiversity Science & Sustainability



CALIFORNIA
ACADEMY OF
SCIENCES

Copyright © 2015 by the California Academy of Sciences

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage or retrieval system, without permission in writing from the publisher.

SCIENTIFIC PUBLICATIONS

Publisher: Margaret Lowman, Ph.D.
Chief, Institute for Biodiversity Science & Sustainability
California Academy of Sciences

EDITORIAL BOARD

Alan E. Leviton, Ph.D., *Editor*
Katherine Piatek, M.A., *Managing Editor*
Michael T. Ghiselin, Ph.D., *Associate Editor*
Tomio Iwamoto, Ph.D., *Associate Editor*
Gary C. Williams, Ph.D., *Associate Editor & Website Coordinator*
Michele L. Aldrich, Ph.D., *Consulting Editor*

COVER IMAGE

Philippine frogs

(See Diesmos, A. et al., *Amphibians of the Philippines, Part I; Checklist of the Species*)
Includes photos by Rafe Brown, Jason Fernandez, and Cameron Siler

COVER DESIGN

Gary C. Williams & Alan E. Leviton
California Academy of Sciences

ISSN 0068–547X

The Proceedings of the California Academy of Sciences is an international journal that accepts manuscripts for publication in the Natural Sciences and selected areas in the Earth Sciences, such as biostratigraphy, regional and global tectonics as they relate to biogeography, and paleoclimatology, and topics in astrobiology, anthropology, as well as the history of science as they relate to institutions of natural history, to individuals, and to activities, such as expeditions and explorations, in the natural sciences.

All manuscripts submitted for publication in any of the Academy's scientific publication series (*Proceedings*, *Occasional Papers*, *Memoirs*) are subject to peer review. Peer review includes both internal and external review, internal review by at least one Academy scientist whose interests parallel those of the submission, and external review, ordinarily by two individuals who are recognized scholars in the field.

Manuscripts accepted for publication are subject to page charges; charges may be waived on a case-by-case basis.

Published by the California Academy of Sciences
55 Music Concourse Drive, Golden Gate Park,
San Francisco, California 94118 U.S.A.

Printed in the United States of America by
Allen Press Inc., Lawrence, Kansas 66044

A New Species of *Nebria* Latreille (Insecta: Coleoptera: Carabidae: Nebriini) from the Spring Mountains of Southern Nevada

David H. Kavanaugh

Department of Entomology, California Academy of Sciences, 55 Music Concourse Drive,
San Francisco, CA 94118, U.S.A. Email: dkavanaugh@calacademy.org

Nebria baumanni Kavanaugh, new species, is described from the Springs Mountains of southern Nevada (type locality: U.S.A., Nevada, Clark County, Spring Mountains, Deer Creek) and is the only known species of the genus endemic to Nevada. It is most closely related to species of the *meanyi* species group based on features of external form and structure and male and female genitalia. This species is named in honor of Dr. Richard Baumann, in recognition of the many contributions by him and his students to our knowledge of arthropod diversity of the Intermountain and Great Basin regions. A key is provided for identification of adults of all *Nebria* species known to occur in Nevada.

KEYWORDS: Coleoptera, Carabidae, Nebriini, *Nebria*, new species, Nevada, Spring Mountains, identification key

Nebria Latreille (1810) is a moderately diverse genus of cool- or cold-adapted carabid beetles restricted to the Holarctic Region and mainly to northern and montane portions of that region. Ledoux and Roux (2005) recorded 384 described species and an additional 158 taxa treated as subspecies in the world's fauna. Since that time, several additional species have been described, including two (Kavanaugh 2008; Kavanaugh and Schoville 2009) from North America. Including the latter, the Nearctic fauna as presently known comprises 85 species, including 31 taxa that previously have been treated as subspecies (Bousquet and Laroche 1993; Bousquet 2012; Kavanaugh 1979, 1981, 1984) but which I now recognize as distinct species.

Recently, I received a few dozen specimens of *Nebria* species from Kipling Will (University of California, Berkeley) for identification. Kip had borrowed most of these specimens from the Monte L. Bean Life Science Museum at Brigham Young University in Provo, Utah, and all were from the state of Nevada. Among those specimens were four from the Spring Mountains of Clark County in southern Nevada that, based on features of external form and structure, were members of the *meanyi* species group. That group currently includes four species: *Nebria giulianii* Kavanaugh (1981), *Nebria lamarckensis* Kavanaugh (1979), *Nebria meanyi* Van Dyke (1925) and *Nebria sylvatica* Kavanaugh (1979). None of these species is known to occur in Nevada, although *N. giulianii* is found in the California portion of the White Mountains, which cross the California-Nevada border and extend a short distance (ca. 15 km) into Esmeralda County, Nevada. So it is certainly possible that *N. giulianii* may occur in suitable habitat (permanent streams or seeps at elevations above 2300 m, if any exist) on the north or northeast slopes of that range, particularly in the vicinity of Boundary Peak.

Although clearly very similar to members of the other *meanyi* group species, the series of four specimens from the Spring Mountains demonstrated several consistent differences from all of them

in external features. Subsequent detailed examination of male and female internal reproductive structures revealed marked differences from members of the previously described species, and provided strong evidence that this series represents a new, undescribed species. The purpose of this paper is to describe this new species of *Nebria*. A key is provided for identification of adult specimens of all *Nebria* species, including this new one, known to occur in the state of Nevada.

MATERIALS AND METHODS

This report is based on study of the four specimens from the Spring Mountains, described here as representing a new species, and tens of thousands of additional specimens representing all previously described Nearctic *Nebria* species. Codons used in the text for collections in which specimens have been deposited include:

BYUC — Monte L. Bean Life Science Museum, Brigham Young University, Provo, UT 84602

CAS — California Academy of Sciences, 55 Music Concourse Drive, San Francisco, CA 94118

Methods used in the present study conform to criteria for ranking taxa as distinct species and techniques for handling specimens as described in Kavanaugh (1979)

Measurements: The only measurement here used is standardized body length (SBL), which equals the sum of the lengths of the head (measured from apex of clypeus to a point on midline at level of posterior margin of compound eye), pronotum (measured from apical margin to basal margin along midline), and elytra (measured along midline from apex of scutellum to apex of the longer elytron).

Dissections: Both male and female genitalia were extracted from specimens relaxed in water immediately after it had boiled and to which a few drops of liquid detergent had been added. Genitalic preparations were then cleared in hot 10% potassium hydroxide solution for five to 10 minutes, each constantly monitored to achieve only a useful degree of clearing. They were then rinsed briefly in 10% acetic acid and then repeatedly in distilled water. Preparations of female genitalic structures were next lightly stained with Chlorazol Black E® (Kodak Corporation). After examination, preparations were stored in glycerin in polyethylene microvials and pinned beneath their specimens of origin.

Illustrations: Digital images of whole specimens and particular structures were taken using a Leica imaging system including an M165C dissecting microscope, DFC550 video camera, and two KL1500 LCD light sources. Stacked images were captured and combined into single montage images using the Leica Application Suite V4.2.0. Plates of images were created using Adobe Photoshop CS5.

SYSTEMATICS

Nebria baumanni Kavanaugh, new species

Figures 1, 2A, 3A, 4A, 5A-B, 6A, 7

TYPE MATERIAL.— Holotype (Fig. 1), a male, deposited in CAS, labeled: “NV Clark Co. Deer Crk Spring Mtns 9 June ’82 Baumann-Clark”/ “HOLOTYPE: *Nebria baumanni* Kavanaugh sp. nov. 2015” [red label]/ “California Academy of Sciences Type No. 18992”. Paratypes (total of 3): 2 females in BYUC and 1 female in CAS with same locality label as holotype, but with the following label: “PARATYPE: *Nebria baumanni* Kavanaugh sp. nov. 2015” [yellow label].




FIGURE 1. Digital images of holotype, *Nebria baumanni* sp. nov. A. Habitus, dorsal aspect; B. Labels for holotype. Scale line = 1.0 mm.

TYPE LOCALITY.— U.S.A., Nevada, Clark County, Spring Mountains, Deer Creek.

DERIVATION OF SPECIES NAME.— It is my great pleasure to name this species in honor of Richard Baumann, who, along with Sean Clark, collected the type series of this species. Throughout his outstanding career, Dr. Baumann and his students have sampled arthropod diversity in the Great Basin and Intermountain regions of the American West more extensively than perhaps any other team. The material that they have collected and made available to others for study, including the specimens described here, have added tremendously to our knowledge of this vast and distinctive region. The species epithet is the Latinized form of his surname name in the genitive case.

DIAGNOSIS.— Adults of this species can be distinguished from those of all other Nearctic *Nebria* species by the following combination of character states: size moderate for genus, SBL of male = 10.4 mm, of females 10.7 to 11.0 mm; body (Fig. 1) black to piceous, dorsal surface without metallic reflection; vertex of head with a pair of paramedial pale spots; elytral microsculpture comprised of moderately impressed isodiametric meshes; antennal scape (Fig. 2A) short, distinctly narrowed basally, widest subapically; pronotum (Fig. 3A) with basal angles rectangular or slightly obtuse, lateral margins with basal sinuation long and deep, sides parallel or slightly convergent basally, with a slight lateral convexity anterior to hind angle in most individuals, lateral

margination ("lateral bead") obliterated from basal one-fifth (posterior to basolateral sinuation), midlateral and basolateral setae present; elytral silhouette subrectangular, distinctly narrowed basally, widest distinctly posterior to middle, lateral margins nearly straight in basal half, humeral angles obtuse and moderately rounded, humeral carinae absent, elytral apices (Fig. 4A) smoothly arcuate laterally and obliquely truncate medially, roundly angulate apically, parascutellar setiferous pore puncture absent; hindwings long but narrowed; median lobe of male aedeagus with shaft thick, evenly arcuate, apical lamella moderately

long, slightly enlarged and evenly rounded apically in left lateral view (Fig. 5A), shaft distinctly thickened in region of apical orifice in dorsal view (Fig. 5B); bursa copulatrix of female genitalia (Fig. 6A) with a posterodorsal lobe and a distinct bursal sclerite in the posterior wall of that lobe, spermathecal duct inserted at base of the bursal sclerite on posterior face of posterodorsal lobe; specimen from Spring Mountains of Clark County, Nevada (Fig. 7).

Members of this species are most similar externally to those of *Nebria giulianii* Kavanaugh and other members of the *meanyi* species group, with which they share similar overall body form and size (Fig. 1). They differ from members of all the other species of this group in the following features: dorsal elytral surface without metallic reflection (distinct metallic blue, green or violet reflection seen in members of all the other species); antennal scape (Fig. 2A) short and markedly narrowed basally (scape longer and not or only slightly narrowed basally (Fig. 2B) in members of all the other species); lateral margination ("lateral bead") of the pronotum obliterated posterior to basal sinuation of lateral margin (lateral margination distinctly defined throughout pronotal length in members of all the other species); elytral apices (Fig. 4A) smoothly arcuate laterally and obliquely truncate medially, roundly angulate apically (broadly and evenly rounded (Fig. 4B) in members of all the other species); median lobe of male aedeagus (Fig. 5A) with shaft thick and apical lamella moderately long and apically slightly enlarged (shaft slender and apical lamella shorter and not apically enlarged (Fig. 5B) in males of all the other species); and bursa copulatrix of female (Fig. 6A) with a posterodorsal lobe and bursal sclerite on posterior face of that lobe (posterodorsal lobe and bursal sclerite absent (Fig. 6B) from females of all the other species).

See the key below to distinguish adults of *N. baumanni* from those of all other species known to occur in Nevada.

SEXUAL DIMORPHISM.—Males and females of this species are similar in size and form and both have two pairs of posterior paramedial setae near the hind margin of sternum VII. The only apparent external difference between the sexes is in the front tarsi: in males the basal three tarsomeres of the protarsi are broad and have dense pads of adhesive setae ventrally, whereas these tarsomeres are slender and without such setal pads in females.




FIGURE 2. Digital images of right antennal scape, dorsal aspect. A. *Nebria baumanni* sp. nov.; B. *Nebria giulianii* Kavanaugh. Scale line = 0.5 mm.

GEOGRAPHICAL DISTRIBUTION.— At present, this species is known only from the upper Deer Creek drainage on the northeastern slope of the Spring Mountains, Clark County, Nevada. It is also the only species known only from the state of Nevada.

HABITAT DISTRIBUTION.— All specimens of the type series were collected under stones along Deer Creek an unspecified distance upstream of a small picnic area just off Deer Creek Road (State Route 158) [Baumann, personal communication].

PHYLOGENETIC RELATIONSHIPS.— Based on characters of external morphology and form and structure of both male and female genitalia, *N. baumanni* is a member of the *meanyi* species group of genus *Nebria* and most closely related to *Nebria giulianii* Kavanaugh (1981) and *Nebria lamarckensis* Kavanaugh (1979).

GEOGRAPHICAL RELATIONS WITH MOST CLOSELY RELATED SPECIES.— The known geographical range of *N. baumanni* is allopatric with respect to the known ranges of all other species of the *meanyi* species group (Fig. 7). *Nebria giulianii* is known

only from the White Mountains in southeastern Mono County, California; and *N. lamarckensis* is known only from the eastern slope of the Sierra Nevada in southwestern Mono County and northwestern Inyo County, California. The nearest localities of these species to the type locality of *N. baumanni* are about 277 and 297 km ENE, respectively, across several lowland gaps of highly inhospitable habitat. The ranges of *N. meanyi* and *N. sylvatica* (see Fig. 7) are far to the north: the former ranging from the mainland coastal mountains of southeastern Alaska and northwestern British Columbia southward along the Cascade Mountain Range from southern British Columbia to Mount Shasta in Siskiyou County, northern California, the latter restricted to the Olympic Mountains of western Washington and Vancouver Island, British Columbia.




FIGURE 3. Digital images of pronotum, dorsal aspect. A. *Nebria baumanni* sp. nov.; B. *Nebria giulianii* Kavanaugh. Scale line = 1.0 mm.





FIGURE 4. Digital images of elytral apices, dorsal aspect; A. *Nebria baumanni* sp. nov.; B *Nebria giulianii* Kavanaugh. Scale line = 0.5 mm.



FIGURES 5. Digital images of median lobe of male genitalia. A-B. *Nebria baumanni* sp. nov.; C-D. *Nebria giulianii* Kavanaugh; A and C. dorsal aspect; B and D. left lateral aspect. Scale line = 1.0 mm.




FIGURE 6. Digital images of female reproductive tract, dorsal aspect. A. *Nebria baumanni* sp. nov.; B. *Nebria giulianii* Kavanagh; badl = anterodorsal lobe of bursa copulatrix; bpdL = posterodorsal lobe of bursa copulatrix; bs = bursal sclerite (in posterior wall of posterodorsal lobe); co = common oviduct; sd = spermathecal duct; sp = spermathecal reservoir; white circle = insertion point of spermathecal duct (at base of posterior face of dorsal lobe). Scale line = 0.5mm.

A Key for Identification of Adults of *Nebria* Species Known to Occur in Nevada

- 1 Head without pale spots on the vertex (to date recorded only from Elko County, Ruby Mountains) *Nebria castanipes* (Kirby)
- 1' Head with a pair of pale reddish spots on the vertex 2
- 2(1') Pronotum with basolateral setae present but midlateral setae absent. 3
- 2' Pronotum with both midlateral and basolateral setae present. 5
- 3(2) Leg color black to rufopiceous (to date recorded only from Elko County, Ruby Mountains) *Nebria edwardsi* Kavanagh
- 3' Leg color pale yellow or tan-orange 4
- 4(3') Pronotum with lateral margin not or only faintly sinuate anterior to obtuse hind angle, lateral explanation moderately broad (widespread in state). *Nebria oblique* LeConte
- 4' Pronotum with lateral margin slightly but distinctly sinuate anterior to rectangular hind angle, lateral explanation moderately narrow (to date recorded only from Washoe County) *Nebria eschscholtzii* Ménériés
- 5(2') Elytral silhouette distinctly ovoid; antennal scape short, thick, ovoid, symmetrical or more convex anteriorly, not narrowed basally; (to date recorded only from Washoe County, Mount Rosa area) *Nebria ovipennis* LeConte
- 5' Elytral silhouette subrectangular or subovoid; antennal scape short to moderately long, varied in shape 6

- 6(5') Antennal scape moderately long, oval and slightly more convex anteriorly, not narrowed basally; elytral silhouette subrectangular; body small to medium-size, SBL males 8.9 to 10.4 mm, females 8.6 to 10.9 mm (to date recorded only from Washoe County) *Nebria rathvoni* LeConte
- 6' Antennal scape short or moderately long, slightly to markedly narrowed basally; elytral silhouette subrectangular or subovoid, body medium- to large-sized, SBL males 10.4 to 12.2 mm, females 10.6 to 13.4 mm 7
- 7(6') Elytral silhouette subovoid, lateral margins more or less arcuate throughout, widest at or slightly posterior to middle; antennal scape moderately long, slightly to moderately narrowed basally; body medium- to large-sized, SBL males 10.5 to 12.2 mm, females 10.6 to 13.4 mm (to date recorded only from Elko County, Jarbridge and Ruby Mountains) *Nebria trifaria* LeConte
- 7' Elytral silhouette (Fig. 1) subrectangular, narrowed and with nearly straight lateral margins in basal half, widest distinctly posterior to middle; antennal scape (Fig. 2A) short, thick apically and markedly narrowed basally; body medium-sized, SBL male 10.4, female 10.7 to 11.0 mm (to date recorded only from Clark County, Spring Mountains) *Nebria baumanni* Kavanaugh, sp. nov.

COMMENTS

I have spent the last four and a half decades collecting and studying *Nebria* in western North America. During that period, I have had the opportunity to visit repeatedly many montane sites throughout the region. At every one of these sites, I’ve recorded the same trend — an upward retreat of the lower limit of the altitudinal range of the species that occur there. The magnitude of this retreat is monumental — an average of about 300 meters for essentially every species examined (unpublished data, manuscript in preparation). Because members of all North American *Nebria* species of which I am aware are cold- or at least cool-adapted general predators, I cannot account for such a region-wide and multispecies phenomenon in any way except for climate warming during the past 40+ years.

The beetles described here as a new species were collected just over 33 years ago, and it will be interesting to see how high one must go to find them now, if their habitat exists there at all. I hope that they can still be found extant in the Spring Mountains, and that documenting the existence of this endemic species for the first time will stimulate biologists in the region to look for them.

ACKNOWLEDGEMENTS

I thank Kipling W. Will (University of California, Berkeley) for making the specimens described herein available to me for study from among material that he had borrowed from the Monte L. Bean Life Science Museum, Brigham Young University. I thank Sean Clark and Richard Baumann at BYU for approving that loan and also for their notes and recollections concerning the collecting of the type series back in 1982. They also approved the deposition of the holotype and one paratype in the collection at CAS. I also want to thank two reviewers, Terry Erwin and Michele Aldrich, for their helpful comments.

FIGURE 7 (right). Map illustrating known localities for members of the *meanyi* species group of *Nebria*; black diamond = *N. baumanni* sp. nov.; black square = *N. giulianii* Kavanaugh; black traingle = *N. lamarckensis* Kavanaugh; black dot = *N. meanyi* Van Dyke; inverted black triangle = *N. sylvatica* Kavanaugh. Scale line = 300 km. Based on “USA Region West landcover location map” from Wikimedia Commons, available at https://upload.wikimedia.org/wikipedia/commons/2/2e/USA_Region_West_landcover_location_map.jpg.



REFERENCES

- BOUSQUET, Y., AND A. LAROCHELLE. 1993. Catalogue of the Geadephaga (Coleoptera: Trachypachidae, Rhyssodidae, Carabidae including Cicindelini) of America north of Mexico. *Memoirs of the Canadian Entomological Society* 167:1–397.
- BOUSQUET, Y. 2012. Catalogue of Geadephaga (Coleoptera, Adephaga) of America, north of Mexico. *ZooKeys* 245:1–1722.
- KAVANAUGH, D.H. 1979. Studies on the Nebriini (Coleoptera: Carabidae), III. New Nearctic *Nebria* species and subspecies, nomenclatural notes, and lectotype designations. *Proceedings of the California Academy of Sciences*, ser. 4, 42:87–133.
- KAVANAUGH, D.H. 1981. Studies on the Nebriini (Coleoptera: Carabidae), IV. Four new *Nebria* taxa from western North America. *Proceedings of the California Academy of Sciences*, ser. 4, 42:435–442.
- KAVANAUGH, D. H. 1984. Studies on the Nebriini (Coleoptera: Carabidae), V. New Nearctic *Nebria* taxa and changes in nomenclature. *Proceedings of the California Academy of Sciences*, ser. 4, 43:159–177.
- KAVANAUGH, D. H. 2008. A new species of *Nebria* Latreille (Insecta: Coleoptera: Carabidae: Nebriini) from the Grand Canyon, Arizona. *Annals of the Carnegie Museum* 77:1–5.
- KAVANAUGH, D. H., and S. D. Schoville. 2009. A new and endemic species of *Nebria* Latreille (Insecta: Coleoptera: Carabidae: Nebriini), threatened by climate change in the Trinity Alps of northern California. *Proceedings of the California Academy of Sciences*, ser. 4, 60:73–84.
- KIRBY, W. 1837. Part 4. The insects. [xxxix + 325 pp.] In: J. Richardson, *Fauna Boreali-Americana; or the Zoology of the Northern Parts of British America*. Josiah Fletcher, Norwich and London, UK.
- LATREILLE, P. A. 1810. *Considérations générale sur l'ordre naturel des animaux composant les classes des Crustacés, des Arachnides, et des Insectes; avec un tableau méthodique de leurs genres, disposés enfamilles*. F. Schoell, Paris, France. 444 pp.
- LECONTE, J. L. 1853. Notes on the classification of the Carabidae of the United States. *Transactions of the American Philosophical Society* (new series) 10:363–403.
- LECONTE, J. L. 1866. Additions to the coleopterous fauna of the United States. I. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1866:361–394.
- LECONTE, J. L. 1878. The Coleoptera of the alpine regions of the Rocky Mountains. *Bulletin of the United States Geological and Geographical Survey of the Territories* 4:447–480.
- LEDoux, G., AND P. ROUX. 2005. *Nebria* (Coleoptera, Nebriidae). *Faune mondiale*. Société Linnéenne de Lyon, Lyon, France. 975 pp.
- MÉNÉTRIÉS, M. 1844. Sur un envoi d'Insectes de la côte N. O. d'Amérique. *Bulletin de la Classe Physico-Mathématique de l'Académie Impériale des Sciences de Saint-Petersbourg* 2:50–63.
- VAN DYKE, E. C. 1925. Studies of western North American Carabinae with descriptions of new species. *The Pan-Pacific Entomologist* 1:111–125.

***Anisotes tablensis* (Acanthaceae), a New Species
from Southwestern Madagascar**

Thomas F. Daniel

*Department of Botany, Institute for Biodiversity and Sustainability, California Academy of Sciences, 55
Music Concourse Drive, San Francisco, California 94118, U.S.A.; Email: tdaniel@calacademy.org*

Anisotes tablensis is described as a new species from an arid region of southwestern Madagascar. It is one of seven species of the genus now known from Madagascar, and it differs from its Malagasy congeners by the following combination of characters: leaves coriaceous to subsucculent, sessile to subsessile, blades 4–23 mm long and 1.4–3 mm wide; bracteoles absent; calyx 6–7 mm long; and corolla externally pubescent with glandular and eglandular trichomes, the lobes of the lower lip 2.5–3.5 mm long. Some of these characters are shared with species from arid regions of southern Madagascar, whereas others are shared with species from less arid regions of northern Madagascar. The conservation status of this apparently rare species is assessed provisionally as Data Deficient.

KEYWORDS: *Anisotes*, Madagascar, endemic species, conservation, pollen, floristics

Since Baden's (1981) revision of it, the Paleotropical genus *Anisotes* Nees has been the subject of several recent studies in Africa (Vollesen 2010, 2015), Madagascar (Daniel et al. 2007, 2013), and the Comoros Archipelago (Daniel 2014). Thirty species are currently recognized in the genus; they occur in mainland Africa (21), the Arabian Peninsula (1), Socotra (1), Madagascar (6), and the Comoros Archipelago (2). All six of the Malagasy species are endemic to that island nation, and most of them appear to be local in occurrence. A seventh locally endemic species is herewith added to the known flora of Madagascar. Although it is from an arid region in southern Madagascar, and shows significant affinities to the two other species of *Anisotes* known from nearby regions, the new species also shows some features in common with congeners from less arid regions in the northern part of the country.

***Anisotes tablensis* T.F. Daniel, sp. nov.**

TYPE. MADAGASCAR: **Toliara:** La Table, ca. 20 km N [*sic*] of Toliara, slope and along ridge, 23°25'26"S, 043°46'03"E, 50–120 m, 19 May 2004 (flr), Z. Rogers et al. 483 (holotype: K!; isotype: MO!). Figures 1, 3.

Divaricately branched shrubs to 4.5 dm tall. Young stems hexagonal, evenly and densely pubescent with a whitish, felt-like covering of antrorsely appressed eglandular trichomes to 0.1 mm long, epidermis not visible, trichomes soon \pm restricted to troughs of internodes. Leaves sessile to subsessile, petioles (if present) to 2 mm long, coriaceous, blades oblong to oblanceolate, 4–23 mm long, 1.4–3 mm wide, 2.9–10.5 \times longer than wide, gradually attenuate at base, rounded at apex, surfaces pubescent like young stems, margin flat. Inflorescence of axillary or terminal sessile short spikes (usually only 2 fertile bracts per spike) to 9 mm long (excluding corollas), distal




FIGURE 1. *Anisotes tablensis* (Rogers et al. 483). A. Habit. B. Leaf. C. Node with axillary inflorescence bearing a flower. D. Distal portion of stamen with anther. Drawn by Tom Davis.

bracts fertile, elliptic, 4.5–5 mm long, 3.6–4 mm wide, apex rounded to obtuse-truncate, abaxial surface puberulent with antrorsely appressed eglandular trichomes to 0.1 mm long and erect trichomes < 0.05 mm long, margin \pm hyaline, proximal bracts sterile and smaller than distal ones. Bracteoles absent. Calyx 5-lobed, 6–7 mm long, lobes oblong-lanceolate, 5–6 mm long, 1–1.2 mm wide, abaxial surface pubescent like bracts (or with antrorsely-appressed eglandular trichomes few or absent). Corolla pinkish red, 27–35 mm long, externally pubescent with erect glandular and flexuose to retrorse eglandular trichomes 0.1–0.2 mm long, tube 12–15 mm long, corolla tube length: corolla length = 0.44–0.49, upper lip 17–21 mm long, lower lip 18–19 mm long with lobes 2.5–3.5 mm long. Stamens 20–23 mm long, filaments pubescent proximally with erect glandular trichomes 0.1–0.2 mm long, glabrous distally, thecae 2–2.3 mm long, glabrous, pollen 3-colporate, 6-pseudocolpate (the two pseudocolpi in one or more mesocolpia sometimes fused near one or both poles), 39–41 μ m polar diameter (P), 26–30 μ m equatorial diameter (E), P:E = 1.37–1.50. Style 28–32 mm long, glabrous, stigma equally 2-lobed, lobes 0.2 mm long. Capsule not seen.

PHENOLOGY.—Flowering: May.

DISTRIBUTION AND HABITAT.—Endemic to southwestern Madagascar (Toliara) where plants occur in the arid thornscrub on soils derived from decomposed coral (sometimes referred to as “coral rag scrub”) at an elevation between 50 and 120 meters (Fig. 2).

CONSERVATION ASSESSMENT.—Because *A. tablensis* is known from a single, relatively recent collection from a population that was not observed during this study, it is difficult to assess its conservation status according to IUCN (2014) guidelines. Based on IUCN criteria for the threatened categories, *A. tablensis* likely has both EOO and AOO (criteria B1 and B2) sufficient for Critically Endangered status, but only one (i.e., one known location; i.e., criterion Ba) of two conditions required for assessment in that category. The only known collection does not occur in a locality that is protected. Lacking further data, this apparently rare species must be provisionally assessed as Data Deficient.

Anisotes tablensis is known only from the type collection. It is the third species of the genus known from the dry regions of southwestern Madagascar. The geographic coordinates given on the



FIGURE 2. La Table in southwestern Madagascar (looking westward toward Mozambique Channel), habitat of *Anisotes tablensis*. Photo by the author.

label of the type are about 2 km southwest of the prominent local mesa-like physiographic feature known as La Table (Fig. 2), from which the epithet is derived. La Table (23°24'32.40"S, 43°46'51.73"E) is a well-known collecting locale in a region of Tertiary limestone about 12 km southeast of the city of Toliara (Tulear) and home to a large number of Acanthaceae. It is also a popular birdwatching site where several rare birds, including the Red-shouldered Vanga (*Calicalicus rufocarpalis*) and Verreaux's Coua (*Coua verreauxi*), are often observed. Given its accessibility, it is surprising that undescribed species are still being found on its slopes and ridge. Two other species of *Anisotes*, *A. divaricatus* T. F. Daniel, Mbola, Almeda & Phillipson and *A. madagascariensis* R. Ben., occur in nearby regions (Daniel et al. 2007), but neither is known from the vicinity of La Table. These three species can be distinguished by the following key:

- 1a. Leaves sessile to subsessile, petioles (if present) to 2 mm long, blades 1.4–3 mm wide, length:width = 2.9–10.5; calyx 6–7 mm long; corolla externally pubescent with glandular and eglandular trichomes throughout, lobes of lower lip 2.5–3.5 mm long. *A. tablensis*
- 1b. Leaves petiolate, petioles to 12 mm long, blades 4.5–38 mm wide, length:width = 0.7–2.7; calyx 1.3–3.5 mm long; corolla externally glabrous or occasionally with a few eglandular trichomes proximally, lobes of lower lip 8–14 mm long 2
- 2a. Leaf blades broadly ovate to ovate-elliptic to elliptic to broadly elliptic, 14–45 mm long; margin of calyx lobes ± densely ciliate; corolla with internal surface conspicuously lighter in color than external surface, corolla tube 9–15 mm long, corolla tube:corolla = 0.23–0.38, upper lip 20–35 mm long and distally whitish to pinkish along margin, lower lip spirally coiled, 18–28 mm long, lobes 12–14 mm long; stamens 26–33 mm long; capsule pubescent with flexuose to antrorse eglandular trichomes 0.05–0.1 mm long *A. madagascariensis*
- 2b. Leaf blades broadly obovate to subcircular to obcordate to obdeltate to oblate, 5–14 mm long; margin of calyx lobes eciliate to sparsely ciliate; corolla with internal surface not conspicuously lighter in color than external surface, corolla tube 14–20 mm long, corolla tube:corolla = 0.46–0.56, upper lip (11–) 14–18 mm long and lacking a pale margin distally, lower lip recurved to reflexed (not spirally coiled), 11–19 mm long, lobes 8–12 mm long; stamens 15–18.5 mm long; capsule ± rugose-granulate but lacking noticeable eglandular trichomes . . . *A. divaricatus*

All three of the species of *Anisotes* from southern Madagascar lack bracteoles and have coriaceous to subsucculent leaves, bracts lacking reticulate venation (or with venation not evident), and similar 3-aperturate pollen. They would appear to pertain to Baden's section *Spiciflori* (Baden 1981; Daniel et al. 2007). Pollen of *A. tablensis* (Fig. 3) greatly resembles that of both *A. madagascariensis* and *A. divaricatus* from southern Madagascar, except for its somewhat smaller polar diameter (39–41 vs. 62.5–7 µm). The four species from northern Madagascar (*A. hygrosopicus* T. F. Daniel, R. Letsara & Martín-Bravo, *A. perplexus* T. F. Daniel, R. Letsara & Martín-Bravo, *A. subcoriaceus* T. F. Daniel, R. Letsara & Martín-Bravo, and *A. venosus* T. F. Daniel, R. Letsara & Martín-Bravo; Daniel et al. 2013) show greater diversity in pollen shape and sculpturing (varying from 2- to 3-aperturate) and have membranous to subcoriaceous leaves, bracteoles, and bracts with reticulate venation evident. *Anisotes tablensis* shows greater similarities to those four species than to the other two southern ones in the following characteristics: calyx length, external pubescence of the corolla, and length of lobes of the lower lip.




FIGURE 3. Pollen of *Anisotes tablensis* (Rogers *et al.* 483). A. Apertural view. B. Interapertural view.

ACKNOWLEDGMENTS

I thank the curators of CAS, G, K, MO, P, and TAN for providing access to specimens; Scott Serata for assistance with scanning electron microscopy; and Rokiman Letsara, E. Magnaghi, and T. Randriambololona for assistance with field work in Madagascar. Field studies were facilitated by the Madagascar Biodiversity Center of the California Academy of Sciences in Antananarivo. Tom Davis skillfully rendered the illustration of *A. tablensis* during a 2015 internship in biological illustration at the California Academy of Sciences. And last but not least, I thank external reviewers C. Anderson and C. Kiel for their helpful comments.

LITERATURE CITED

- BADEN, C. 1981. The genus *Anisotes* (Acanthaceae), a taxonomic revision. *Nordic Journal of Botany* 1:623–664.
- DANIEL, T. F. 2014. Taxonomy of *Anisotes* Nees (Acanthaceae: Justicieae) in the Comoros Archipelago and a preliminary list of Acanthaceae from the islands. *Candollea* 69:45–54.
- DANIEL, T. F., R. LETSARA, AND S. MARTÍN-BRAVO. 2013. Four new species of *Anisotes* (Acanthaceae) in Madagascar. *Novon* 22:396–408.
- DANIEL, T. F., B. A. V. MBOLA, F. ALMEDA, AND P. B. PHILLIPSON. 2007. *Anisotes* (Acanthaceae) in Madagascar. *Proceedings of the California Academy of Sciences*, ser. 4, 58:121–131.
- IUCN. 2014. Guidelines for Using the IUCN Red List Categories and Criteria, Version 11. Standards and Petitions Subcommittee of the IUCN Species Survival Commission. <<http://www.iucnredlist.org/documents/RedListGuidelines.pdf>> [accessed 16 November 2015].
- VOLLESEN, K. 2010. *Anisotes*. Pages 651–663 in H. J. Beentje, ed., *Flora of Tropical East Africa, Acanthaceae*, part 2. Royal Botanic Gardens, Kew, England.
- VOLLESEN, K. 2015. *Anisotes*. Page 236 in J. R. Timberlake and E. S. Martins, eds., *Flora Zambesiaca*, vol. 8(6). Royal Botanic Gardens, Kew, England.

Page intentionally left blank

Amphibians of the Philippines, Part I: Checklist of the Species

Arvin C. Diesmos^{1,†}, **Jessa L. Watters**^{2,†}, **Nicholas A. Huron**^{2,3}, **Drew R. Davis**⁴,
Angel C. Alcala⁵, **Ronald I. Crombie**⁶, **Leticia E. Afuang**⁷, **Genevieve Gee-Das**⁸,
Rogelio V. Sison^{9,§}, **Marites B. Sanguila**¹⁰, **Michelle L. Penrod**², **Marie J. Labonte**²,
Conner S. Davey², **E. Austin Leone**², **Mae L. Diesmos**¹¹, **Emerson Y. Sy**¹²,
Luke J. Welton¹³, **Rafe M. Brown**¹³, and **Cameron D. Siler**^{2,3,*}

¹ Herpetology Section, Zoology Division, Philippine National Museum, Rizal Park, Burgos St., Manila, Philippines; ² Sam Noble Oklahoma Museum of Natural History, University of Oklahoma, 2401 Chautauqua Ave., Norman, OK 73072-7029, USA; ³ Department of Biology, University of Oklahoma, 730 Van Vleet Oval, Norman, OK 73019, USA; ⁴ Department of Biology, University of South Dakota, 414 East Clark St., Vermillion, SD 57069, USA; ⁵ Angelo King Center for Research and Environmental Management, Silliman University, 2F Marine Laboratory, Bantayan, Dumaguete City, Philippines; ⁶ 1489 Tunnel Rd, Santa Barbara, CA 93105, USA; ⁷ Animal Biology Division, Institute of Biological Sciences, College of Arts and Sciences, University of the Philippines Los Baños, Laguna 4031, Philippines; ⁸ Herpetofauna Research Group, Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia; ⁹ Herpetology Section, Zoology Division, National Museum of the Philippines, Padre Burgos Avenue, Ermita, 1000 Manila, Philippines; ¹⁰ Father Saturnino Urios University, 8600 Butuan City, Philippines; ¹¹ University of Santo Tomas, Manila, Philippines; ¹² Philippine Center for Terrestrial and Aquatic Research, 1198 Benavidez Street, Unit 1202, Sta. Cruz, Manila, Philippines; ¹³ Biodiversity Institute and Department of Ecology and Evolutionary Biology, University of Kansas, 1345 Jayhawk Blvd., Lawrence, KS 66045-7593, USA

The herpetological fauna of the Philippine Islands is high in diversity and endemism (Brown and Diesmos 2009; Brown et al. 2013; Diesmos et al. 2014), yet faces threats such as habitat modification and loss, natural catastrophes (i.e., Typhoon Haiyan), invasive species, hunting for food or the pet trade, and the spread of chytrid fungus (Sodhi et al. 2004; Diesmos et al. 2006, 2012; Rowley et al. 2010; Brown et al. 2012). New species descriptions have been steadily rising since the early 1990s due to increased sampling, an awareness of species boundaries based on phylogenetic studies, and changes in our understanding of what defines a species (Figure 1 [p. 489]; Brown et al. 2001, 2008, 2013; Diesmos et al. 2002, 2012; Diesmos and Brown 2011; Brown and Stuart 2012). Developing a complete species list for amphibians is essential for conservation planning and informed management decisions. Previous lists (Brown 2007; Diesmos and Brown 2011; Diesmos et al. 2014) were derived in part from working compendiums, developed and distributed separately by RIC and ACD; these simple lists focused on taxonomic and conservation status of the included species, respectively, but were of limited use for other purposes.

Herein we provide a comprehensive checklist of Philippine amphibian diversity, created by searching worldwide museum databases for Philippines taxa, augmented with a thorough review of recently published new species descriptions. Museum records from 33 museums were obtained either through direct contact with museum websites and personnel or through database portals such

†Authors made equal contributions to this manuscript

§Deceased

*Corresponding author, e-mail: camsiler@ou.edu

as VertNET (<<http://vertnet.org>>) and Global Biodiversity Information Facility (GBIF, <<http://www.gbif.org>>). In total, we reviewed 43,222 specimen records.

In this checklist, information associated with each species has the following sequence: (1) the original source, (2) a non-exhaustive, representative synonymy (see also comments below), (3) the type locality as reported in the authoritative description and holotype catalog number, if known, and (4) distribution within the Philippines and identification of endemism. Distributions are given as island names only, with species considered present on each reported island. Full citations for all authoritative descriptions are provided in the Literature Cited section.

Distribution dot maps were created based on a total of 4,015 unique localities from the georeferenced museum records described above using ArcMap v.10.3.1 (Figures 2–30, 43). Due to lack of precise locality data or coordinates for some museum records, a proportion of known species occurrences may not have been included in the dot maps. Topographic maps were created in ArcMap v.10.3.1 using the digital elevation model (DEM) layers based on NASA's Shuttle Radar Topographic Mission (SRTM). The SRTM data are available for free at approximately 90 meters resolution (3 arc-second projections; Reuter et al. 2007; CIAT-CSI SRTM 2015). Representative photos of most currently recognized species have also been provided (Figures 31–42, 44).

Although every effort was made to provide detailed taxon-specific synonyms for every species, the emphasis of this study was to provide an updated documentation and concise overview of the current diversity of amphibians in the Philippines. We direct readers to additional primary sources and amphibian biodiversity information products for original species descriptions and complete synonymies: Amphibian Species of the World (<<http://research.amnh.org/vz/herpetology/amphibia/>>; Frost 2015) and AmphibiaWeb (<<http://amphibiaweb.org>>; Amphibia Web 2015).

Museum designation symbolic codes follow Sabaj Pérez (2014): British Museum of Natural History (BMNH); California Academy of Sciences (CAS); California Academy of Sciences, Stanford University (CAS-SU); Carnegie Museum (CM); Field Museum of Natural History (FMNH); Museum of Comparative Zoology, Harvard University (MCZ); Museum National d'Histoire Naturelle (MNHN); Museo Civico di Storia Naturale (MSNG); Naturhistorisches Museum (NHMW of NMW); National Museum of the Philippines (formerly Philippine National Museum; NMPH or PNM); Naturalis Biodiversity Center (RMNH); Senckenberg Forschungsinstitut und Naturmuseum (SMF); National Museum of Natural History, Smithsonian Institution (USNM); Museum für Naturkunde (ZMB).

Amphibians of the Philippines, Part II, which will include identification keys for the amphibian fauna, will be issued in 2016.

Class Amphibia, Order Anura
Family Bombinatoridae

***Barbourula busuangensis* Taylor and Noble 1924**

Barbourula busuangensis Taylor and Noble 1924

Type locality and holotype specimen: Philippines, “small stream in the southern part of Busuanga [Island], the largest island of the Calamianes group” (Taylor and Noble 1924) (MCZ 14004).

Philippine distribution: Balabac, Busuanga, Culion, Palawan (Endemic).

Figures 3 and 31.

Family Bufonidae

***Ansonia mcgregori* (Taylor 1922)**

Bufo mcgregori Taylor 1922, 1944; Slevin and Leviton 1956; Van Tuijl 1995

Ansonia muelleri Inger 1954

Ansonia mcgregori, Inger 1960

Type locality and holotype specimen: Philippines, Mindanao, Zamboanga, near Pasonanka (= Pasonanca) (CAS 61839).

Philippine distribution: Mindanao (Endemic).

Figures 3 and 31.

***Ansonia muelleri* (Boulenger 1887)**

Bufo muelleri Boulenger 1887; Frost 1985

Ansonia muelleri, Inger 1954; Alcalá and Brown 1998

Type locality and holotype specimen: Philippines, Mindanao (BMNH 1947.2.20.57).

Philippine distribution: Dinagat, Mindanao (Endemic).

Figures 3 and 31.

***Ingerophrynus philippinicus* (Boulenger 1887)**

Bufo philippinicus Boulenger 1887; Iskandar 1998; Inger, 1999

Bufo divergens Mocquard 1890

Bufo biporcatus philippinicus, Inger 1954; Alcalá and Brown 1998

Ingerophrynus philippinicus, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Type locality and holotype specimen: Philippines, Palawan, Puerto Princesa (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Balabac, Busuanga, Culion, Dumarán, Palawan (Endemic).

Figures 3 and 31.

***Pelophryne albotaeniata* Barbour 1938**

Pelophryne albotaeniata Barbour 1938

Pelophryne albotaeniata albotaeniata, Inger 1954

Type locality and holotype specimen: Philippines, Palawan, Thumb Peak (4,500 ft. elev.) (MCZ 23291).

Philippine distribution: Palawan (Endemic).

Figure 4.

Pelophryne brevipes* (Peters 1867)Hylaplesia brevipes* Peters 1867*Bufo brevipes*, Cope 1867; Boulenger 1882*Pelophryne brevipes*, Barbour 1938

Type locality and holotype specimen: Philippines, Mindanao, Zamboanga (NMW 16554, syntype).

Philippine distribution: Basilan, Mindanao (Endemic).

Figures 4 and 31.

Pelophryne lighti* (Taylor 1920)Nectophryne lighti* Taylor 1920*Pelophryne lighti*, Barbour 1938; Inger 1960*Pelophryne brevipes*, Inger 1954*Pelophryne albotaeniata lighti*, Inger 1954

Type locality and holotype specimen: Philippines, Mindanao, Agusan Province, Bunawan (E. H. Taylor Collection No. 189, destroyed).

Philippine distribution: Bohol, Leyte, Mindanao, Samar (Endemic).

Figures 4 and 31.

Rhinella marina* (Linnaeus 1758)Rana marina* Linnaeus 1758*Bufo brasiliensis* Laurenti 1768*Rana gigas* Walbaum 1784*Rana humeris-armata* Lacépède 1788*Rana humeris-armata*, Bonnaterre 1789*Bufo marinus* Schneider 1799; Gravenhorst 1829*Bufo agua* Sonnini de Manoncourt and Latreille 1801*Rana brasiliensis* Shaw 1802*Bufo horridus* Daudin 1802*Bufo humeralis* Daudin 1803*Bombinator maculatus* Merrem 1820*Rana maxima* Merrem 1820*Bombinator horridus* Merrem 1820*Bufo maculiventris* Spix 1824*Bufo lazarus* Spix 1824*Bufo albicans* Spix 1824*Bufo horribilis* Weigmann 1833*Docidophryne agua* Fitzinger 1843*Docidophryne Lazarus* Fitzinger 1861*Phrynoidis agua* Cope 1862*Bufo marinus* var. *horribilis* Peters 1873; Lynch and Fugler 1965*Bufo marinus* var. *fluminensis* Jiménez de la Espada 1875*Bufo marinus* var. *napensis* Jiménez de la Espada 1875*Bufo pithecodactylus* Werner 1899*Bufo marinus*, Barbour and Noble 1920*Bufo marinus marinus*, Schmidt 1932*Bufo angustipes* Taylor and Smith 1945*Bufo pythecodactylus* Rivero 1961

Chaunus marinus, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Rhinella marina, Chaparro, Pramuk, and Gluesenkamp 2007

Rhinella marinus, Pramuk, Robertson, Sites, and Noonan 2008

Type locality and holotype specimen: America (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Alabat, Bohol, Calayan, Catanduanes, Cebu, Cocomo, Dinagat, Gigantes Norte, Leyte, Lubang, Luzon, Marinduque, Masbate, Mindanao, Mindoro, Negros, Palawan, Panay, Polillo, Romblon Island Group, Sicogon, Samar, Ticao, Verde (Introduced; Diesmos et al. 2006).

Figures 4 and 31.

Family Ceratobatrachidae

Alcalus mariae (Inger 1954)

Micrixalus mariae Inger 1954

Ingerana (*Ingerana*) *mariae*, Dubois 1987 “1986”

Taylorana mariae, Fei, Ye, and Jiang 2010

Alcalus mariae, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Palawan, Mantalingajan Range, south slope of Mount Balabag (FMNH 51360).

Philippine distribution: Palawan (Endemic).

Figures 12 and 35.

Platymantis banahao Brown, Alcalá, Diesmos, and Alcalá 1997

Platymantis banahao Brown, Alcalá, Diesmos, and Alcalá 1997

Platymantis (*Lahatnanguri*) *banahao*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Quezon Province, NE slope of Mt. Banahao (1,100 m elev.) (CAS 201208).

Philippine distribution: Luzon (Endemic).

Figures 5 and 32.

Platymantis bayani Siler, Alcalá, Diesmos, and Brown 2009

Platymantis bayani Siler, Alcalá, Diesmos, and Brown 2009

Platymantis (*Tahananpuno*) *banahao*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Samar, Eastern Samar Province, Municipality of Taft, Barangay San Rafael, Taft Forest (11.80255°N, 125.29276°E; WGS84; 140 m elev.) (PNM 9501).

Philippine distribution: Samar (Endemic).

Figures 5 and 32.

Platymantis biak Siler, Diesmos, Likem, Diesmos, and Brown 2010

Platymantis biak Siler, Diesmos, Linkem, Diesmos, and Brown 2010

Platymantis (*Lahatnanguri*) *biak*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Bulacan Province, Municipality of San

Miguel and Doña Remedios Trinidad, Barangay Biak na Bato (15.1084°N, 121.0724°E; 190 m elev.) (PNM 9679).

Philippine distribution: Luzon (Endemic).

Figure 5.

***Platymantis cagayanensis* Brown, Alcala, and Diesmos 1999**

Platymantis cagayanensis Brown, Alcala, and Diesmos 1999

Platymantis (Lupacolus) cagayanensis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Cagayan Province, Central Cordillera, Santa Praxedes Town, Taggat Forest Reserve (50–100 m elev.) (PNM 6691).

Philippine distribution: Luzon, Palau (Endemic).

Figures 5 and 32.

***Platymantis cornutus* (Taylor 1922)**

Cornufer cornutus Taylor 1922

Platymantis cornutus, Zweifel 1967

Platymantis cornuta, Günther 1999

Platymantis (Lahatnanguri) cornutus, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Mountain Province, Kalinga, Balbalan (CAS 61476).

Philippine distribution: Luzon (Endemic).

Figures 6 and 32.

***Platymantis corrugatus* (Duméril 1853)**

Hylodes corrugatus Duméril 1853

Platymantis plicifera Günther 1858

Hylodes (Halophilus) corrugatus, Cope 1862

Halophila (Platymantis) plicifera, Peters 1863

Platymantis corrugata, Boulenger 1918; Günther 1999

Rana (Platymantis) rugata Van Kampen 1923

Platymantis corrugatus corrugatus, Loveridge 1948

Platymantis (Tagomukhus) corrugatus, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Java (in error, according to Inger 1954) (MNHNP 4884).

Philippine distribution: Bohol, Cagraray, Camiguin Sur, Camotes Island Group, Catanduanes, Cebu, Dinagat, Leyte, Luzon, Mindanao, Mindoro, Negros, Panay, Polillo, Ponson, Poro, Rapu-Rapu, Samar, Sicogon, Siquijor (Endemic).

Figures 6 and 32.

***Platymantis diesmosi* Brown and Gonzalez 2007**

Platymantis diesmosi Brown and Gonzalez 2007

Platymantis (Tahananpuno) diesmosi, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Albay Province, Municipality of Tiwi, Barangay Banshaw, Sitio Purok 7, area known locally as ‘Tamagong,’ NW slope of Mt. (13.436667°N, 123.59°E; WGS84; 950 m elev.) (PNM 8499).

Philippine distribution: Luzon (Endemic).
 Figures 6 and 32.

***Platymantis dorsalis* (Duméril 1853)**

Cornufer dorsalis Duméril 1853

Hylodes (Halophilus) dorsalis, Cope 1862

Halophila jagorii Peters 1863

Halophila platydactyla Günther 1864

Platymantis meyeri Günther 1873; Boulenger 1918

Cornufer jagorii, Boulenger 1882

Cornufer meyeri, Boulenger 1882; Inger 1954

Cornufer laticeps Taylor 1920

Rana (Platymantis) dorsalis, Guibé 1950 “1948”

Cornufer dorsalis, Brown and Inger 1964; Brown 1965

Platymantis dorsalis, Zweifel 1967; Brown and Alcala 1970a, b

Platymantis (Lupacolus) dorsalis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Java (in error, according to Barbour 1923) (MNHNP 4880).
 Philippine distribution: Alabat, Calagna-an, Cagraray, Catanduanes, Cebu, Danjugan, Leyte, Lubang, Luzon, Marinduque, Masbate, Negros, Pan de Azucar, Panay, Polillo, Rapu-Rapu, Sicogon, Ticao (Endemic).

Figures 6 and 32.

***Platymantis guentheri* (Boulenger 1882)**

Cornufer guentheri Boulenger 1882; Inger 1954

Cornufer worcesteri Stejneger 1905

Cornufer ingeri Brown and Alcala 1963

Platymantis ingeri, Zweifel 1967

Platymantis guentheri, Zweifel 1967

Platymantis (Tahananpuno) guentheri, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Dinagat (BMNH 1947.2.31–34).

Philippine distribution: Biliran, Bohol, Dinagat, Leyte, Mindanao, Samar (Endemic).

Figures 7 and 33.

***Platymantis hazelae* (Taylor 1920)**

Philautus hazelae Taylor 1920

Cornufer rivularis Taylor 1922

Rhacophorus (Philautus) hazelae, Ahl 1931

Platymantis hazelae, Inger 1954; Zweifel 1967; Brown and Alcala 1970a, b

Cornufer hazelae, Inger 1954; Brown 1965

Platymantis (Tirahanulap) hazelae, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, central northern Negros, Canlaon Volcano (ca. 1,000 m elev.) (CM 3427).

Philippine distribution: Negros, Masbate (Endemic).

Figures 7 and 33.

Platymantis indepressus* Brown, Alcala, and Diesmos 1999Platymantis indepressus* Brown, Alcala, and Diesmos 1999*Platymantis indepressa*, Günther 1999*Platymantis (Lupacolus) indepressus*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Mt. Banahao (1,080 m elev.) (CAS 201196)

Philippine distribution: Luzon (Endemic).

Figure 7.

Platymantis insulatus* Brown and Alcala 1970Platymantis insulatus* Brown and Alcala 1970a, b*Platymantis insulata*, Günther 1999*Platymantis (Lahatnanguri) insulatus*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Gigante Sur Island (CAS 117441).

Philippine distribution: Gigante Norte, Gigante Sur (Endemic).

Figures 7 and 33.

Platymantis isarog* Brown, Brown, Alcala, and Frost 1997Platymantis isarog* Brown, Brown, Alcala, and Frost 1997*Platymantis reticulatus* Brown, Brown, and Alcala 1997*Platymantis (Tirahanulap) isarog*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, southeastern Luzon, Camarines Peninsula, Mt. Isarog (1,200–1,300 m elev.) (CAS 197218).

Philippine distribution: Luzon (Endemic).

Figures 8 and 33.

Platymantis lawtoni* Brown and Alcala 1974Platymantis lawtoni* Brown and Alcala 1974*Platymantis (Tirahanulap) lawtoni*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Tablas, Dubduban (ca. 800 ft. elev.) (CAS 135732)

Philippine distribution: Romblon Island Group (Endemic).

Figures 8 and 33.

Platymantis levigatus* Brown and Alcala 1974Platymantis levigatus* Brown and Alcala 1974*Platymantis levigata*, Günther 1999*Platymantis (Lahatnanguri) levigatus*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Tablas, San Agustin, Dubduban (ca. 650 ft. elev.) (CAS 136097).

Philippine distribution: Romblon Island Group (Endemic).

Figures 8 and 33.

Platymantis luzonensis* Brown, Alcala, Diesmos, and Alcala 1997Cornifer guentheri*, Inger 1954*Platymantis guentheri*, Brown and Alcala 1970*Platymantis luzonensis* Brown, Alcala, Diesmos, and Alcala 1997*Platymantis (Tahananpuno) luzonensis*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Laguna Province, Mt. Makiling (ca. 600 m elev.) (CAS 196368).

Philippine distribution: Luzon, Polillo (Endemic).

Figures 8 and 33.

Platymantis mimulus* Brown, Alcala, and Diesmos 1999Platymantis mimulus* Brown, Alcala, and Diesmos 1997*Platymantis mimula*, Günther 1999*Platymantis (Lupacolus) mimulus*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Laguna Province, Los Banos, Mt. Maquiling (ca. 400 m elev.) (CAS 136097).

Philippine distribution: Luzon (Endemic).

Figure 9.

Platymantis montanus* (Taylor 1922)Cornufer montanus* Taylor 1922*Platymantis montanus*, Zweifel 1967*Platymantis montana*, Günther 1999*Platymantis (Tirahanulap) montanus*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Laguna Province, Mount Banahao (ca. 1,500 m elev.) (CAS 61179).

Philippine distribution: Luzon (Endemic).

Figures 9 and 33.

Platymantis naomii* Alcala, Brown, and Diesmos 1998Platymantis naomii* Alcala, Brown, and Diesmos 1998*Platymantis naomiae*, Iskandar and Colijn 2000*Platymantis (Lupacolus) naomii*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, southeast slope of Mt. Banahao on Tayabas side (ca. 1,400 m elev.) (CAS 204746).

Philippine distribution: Luzon (Endemic).

Figure 9.

Platymantis negrosensis* Brown, Alcala, Diesmos, and Alcala 1997Platymantis negrosensis* Brown, Alcala, Diesmos, and Alcala 1997*Platymantis (Tahananpuno) negrosensis*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Negros, Negros Oriental Province, Lake Balinsasayao (CAS 137416).

Philippine distribution: Negros, Panay (Endemic).

Figures 9 and 34.

***Platymantis paengi* Siler, Linkem, Diesmos, and Alcala 2007**

Platymantis paengi Siler, Linkem, Diesmos, and Alcala 2007

Platymantis (Lupacolus) paengi, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Panay, Antique Province, Municipality of Pandan, Barangay Duyong, in an area known locally as ‘Mt. Lihidan’ (11.41465°N, 122.10465°E; WGS84; 180 m elev.) (PNM 9239).

Philippine distribution: Panay (Endemic).

Figures 10 and 34.

***Platymantis panayensis* Brown, Brown, and Alcala 1997**

Platymantis panayensis Brown, Brown, and Alcala 1997

Platymantis (Tirahanulap) panayensis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Panay, Aklan Province, Libacao, northwest ridge approach to Mt. Madja-as (ca. 1,410 m elev.) (PNM 2495).

Philippine distribution: Panay (Endemic).

Figure 10.

***Platymantis polillensis* (Taylor 1922)**

Philautus polillensis Taylor 1922

Rhacophorus polillensis, Ahl 1931

Cornufer polillensis, Inger 1954

Platymantis polillensis, Zweifel 1967

Platymantis polilloensis, Alcala 1986; Brown, Brown, and Alcala 1997

Platymantis (Tirahanulap) polillensis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Polillo, near the southern end of island (CAS 62250).

Philippine distribution: Luzon, Polillo (Endemic).

Figures 10 and 34.

***Platymantis pseudodorsalis* Brown, Alcala, and Diesmos 1999**

Platymantis pseudodorsalis Brown, Alcala, and Diesmos 1999

Platymantis (Lupacolus) pseudodorsalis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Mt. Banahao (PNM 6689).

Philippine distribution: Luzon (Endemic).

Figure 10.

***Platymantis pygmaeus* Alcala, Brown, and Diesmos 1998**

Platymantis pygmaeus Alcala, Brown, and Diesmos 1998

Platymantis pygmaea, Günther 1999

Platymantis (Lahatnanguri) pygmaeus, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Isabela Province, Municipality of Palanan, Barangay Didian, Sitio Natapdukan, Northern Sierra Madre Natural Park (16.9655°N, 122.4038°E; WGS84; 55–65 m elev.) (PNM 6255).

Philippine distribution: Luzon, Sibuyan (Endemic).
 Figures 11 and 34.

***Platymantis quezoni* Brown, De Layola, Lorenzo, Diesmos, and Diesmos 2015**

Platymantis (Lupacolus) quezoni Brown, De Layola, Lorenzo, Diesmos, and Diesmos 2015

Platymantis “sp. 27”, Brown, Siler, Richards, Diesmos, and Cannatella, 2015

Type locality and holotype specimen: Philippines, Luzon, Quezon Province, Municipality of Atimonan, Barangay Malinao Ilaya, Quezon Protected Landscape (13.989°N, 121.818°E; WGS84; 275 m elev.) (PNM 9817, formerly KU 339542).

Philippine distribution: Luzon (Endemic).
 Figures 43 and 44.

***Platymantis rabori* Brown, Alcala, Diesmos, and Alcala 1997**

Platymantis rabori Brown, Alcala, Diesmos, and Alcala 1997

Platymantis (Tahananpuno) rabori, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Bohol, Sierra Bullones, Cantaub (CAS 136889).
 Philippine distribution: Bohol, Leyte, Mindanao, Samar (Endemic).
 Figures 11 and 34.

***Platymantis sierramadrensis* Brown, Alcala, Ong, and Diesmos 1999**

Platymantis sierramadrensis Brown, Alcala, Ong, and Diesmos 1999

Platymantis (Tirahanulap) sierramadrensis, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Quezon Province, Municipality of General Nakar, Barangay Umiray, at Sitio Mapidjas (PNM 6465).
 Philippine distribution: Luzon (Endemic).
 Figures 11 and 34.

***Platymantis spelaeus* Brown and Alcala 1982**

Platymantis spelaeus Brown and Alcala 1982

Platymantis spelaea, Günther 1999

Platymantis (Lupacolus) spelaeus, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Negros, southern Negros Oriental, Basay, Tiya-banan Barrio, in limestone cave (CAS 153469).
 Philippine distribution: Negros (Endemic).
 Figures 11 and 34.

***Platymantis subterrestris* (Taylor 1922)**

Cornufer subterrestris Taylor 1922; Inger 1954

Platymantis subterrestris, Zweifel 1967; Brown and Alcala 1970

Platymantis (Tirahanulap) subterrestris, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Mountain Province, near kilometer 101 on the Mountain Trail (CAS 61518).
 Philippine distribution: Luzon (Endemic).
 Figures 12 and 34.

Platymantis taylori* (Brown, Alcalá, and Diesmos 1999)Platymantis taylori* Brown, Alcalá, and Diesmos 1999*Platymantis (Lupacolus) taylori*, Brown, Siler, Richards, Diesmos, and Cannatella 2015

Type locality and holotype specimen: Philippines, Luzon, Province of Isabela, Municipality of Palanan, Barangay Didian, eastern Sierra Madre Mountains in Sitio Natapdukan (PNM 6884).

Philippine distribution: Luzon (Endemic).

Figures 12 and 35.

Family Dicroglossidae***Fejervarya moodiei* (Taylor 1920)***Rana cancrivora* Gravenhorst 1829*Rana tigrina angustopalmata* Van Kampen 1907; Barbour 1912*Rana tigrina* var. *cancrivora*, Boulenger 1918*Rana cancrivora*, Annandale 1918*Rana (Rana) crancrivora*, Boulenger 1920*Rana moodiei*, Taylor 1920*Rana cancrivora cancrivora*, Dunn 1928; Inger 1954*Rana cancrivora raja* Smith 1930*Dicroglossus cancrivorus*, Deckert 1938*Rana raja*, Taylor 1962*Rana (Euphlyctis) cancrivora*, Dubois 1981*Euphlyctis cancrivora*, Poynton and Broadley 1985*Limnonectes (Hoplobatrachus) cancrivorus*, Dubois 1987 “1986”*Limnonectes (Hoplobatrachus) moodiei*, Dubois 1987 “1986”*Limnonectes (Hoplobatrachus) raja*, Dubois 1987 “1986”*Limnonectes (Fejervarya) raja*, Dubois 1992*Limnonectes (Fejervarya) cancrivorus*, Dubois 1992*Fejervarya raja*, Iskandar 1998*Fejervarya cancrivora*, Iskandar 1998*Fejervarya moodiei*, Dubois and Ohler 2000

Type locality and holotype specimen: Indonesia, Java (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Apo, Bohol, Boracay, Busuanga, Cagraray, Calagna-an, Caluya, Camiguin Sur, Cebu, Clara, Cuyo, Dumarán, Dinagat, Gigantes Norte, Gigantes Sur, Guimaras, Inampulugan, Jau, Lapinin Chico, Lapinig Grande, Leyte, Lubang, Luzon, Mactán, Marinduque, Masbate, Mindanao, Mindoro, Negros, Pacijan, Palawan, Pan de Azúcar, Panay, Polillo, Ponson, Rapu-Rapu, Romblon Island Group, Sicogon, Sulu Archipelago, Ticao, Tintiman, Verde (Non-endemic).

Figures 12 and 34.

Fejervarya vittigera* (Wiegmann 1834)Rana vittigera* Wiegmann 1834*Rana limnocharis vittigera*, Inger 1954*Rana (Fejervarya) vittigera*, Dubois 1984*Euphlyctis limnocharis vittigera*, Poynton and Broadley 1985*Limnonectes (Fejervarya) vittiger*, Dubois 1987 “1986”

Rana limnocharis, Alcala and Brown 1998

Fejervarya vittigera, Iskandar 1998

Type locality and holotype specimen: Philippines, southern Luzon, Laguna Bay (CAS 61636).

Philippine distribution: Bohol, Cagraray, Caluya, Camiguin Sur, Cebu, Cocomo, Dinagat, Guimaras, Leyte, Lubang, Luzon, Marinduque, Masbate, Mindanao, Mindoro, Negros, Palawan, Pan de Azucar, Panay, Polillo, Romblon Island Group (Endemic).

Figures 13 and 35.

***Hoplobatrachus rugulosus* (Wiegmann 1834)**

Rana chinensis Osbeck 1765

Rana rugulosa Wiegmann 1834; Annandale 1918; Alcala and Brown 1998

Rana tigrina pantherina Steindachner 1867; Boulenger 1920; Taylor and Elbel 1958

Hydrostentor pantherinus, Steindachner 1867

Rana tigrina, Flower 1899

Rana esculenta chinensis, Wolterstorff 1906

Rana burkilli Annandale 1910

Rana tigrina var. *burkilli*, Boulenger 1918

Rana tigrina rugulosa, Smith 1930; Fang and Chang 1931

Rana (Euphlyctis) rugulosa, Dubois 1981

Euphlyctis tigrina rugulosa, Poynton and Broadley 1985

Limnonectes (Hoplobatrachus) rugulosus, Dubois 1987 “1986”

Tigrina rugulosa, Fei, Ye, and Huang 1990

Hoplobatrachus rugulosus, Dubois 1992

Hoplobatrachus chinensis, Ohler, Swan, and Daltry, 2002

Type locality and holotype specimen: China, vicinity of Canton (ZMB 3721).

Philippine distribution: Caluya, Luzon, Masbate, Mindoro, Panay (Introduced; Diesmos et al. 2006).

Figures 13 and 35.

***Limnonectes acanthi* (Taylor 1923)**

Rana macrodon blythii (part) Boulenger 1920

Rana acanthi Taylor 1923; Taylor and Elbel 1958

Rana macrodon acanthi, Inger 1954; Brown and Alcala 1955

Rana magna acanthi, Inger 1958

Limnonectes (Limnonectes) acanthi, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Calamian Islands, Busuanga Island (CAS 32577).

Philippine distribution: Balabac, Busuanga, Culion, Mindoro, Moro, Palawan (Endemic).

Figures 13 and 35.

***Limnonectes diuatus* (Brown and Alcala 1977)**

Rana diuata Brown and Alcala 1977

Limnonectes (Limnonectes) diuatus, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Mindanao, Agusan del Norte Province, Cabadbaran, Diuata Mountains, south side of Mt. Hilong-hilong, Taguibo River (ca. 1,000 m elev.) (CAS 133500).

Philippine distribution: Mindanao (Endemic).

Figure 13.

Limnonectes ferner* Siler, McVay, Diesmos, and Brown 2009Limnonectes ferner* Siler, McVay, Diesmos, and Brown 2009

Type locality and holotype specimen: Philippines, Mindanao, Davao Del Norte Province, Municipality of Monkayo, Mt. Pasian in the Simulaw River Drainage, 2.3 km N, 1.0 km E of peak (7.971183°N, 126.297367°E; WGS84; 1,409 m elev.) (PNM 9506).

Philippine distribution: Mindanao (Endemic).

Figure 14.

Limnonectes leytensis* (Boettger 1893)Hylarana mindanensis* Girard 1853*Rana mindanensis* Boettger 1886*Rana leytensis*, Boettger 1893; Inger 1966*Rana microdisca* Boulenger 1920*Rana microdisca leytensis*, Inger 1954; Mertens 1967*Limnonectes* (*Limnonectes*) *leytensis*, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Mindanao, unknown caldera (SMF 4931).

Philippine distribution: Basilan, Bohol, Camiguin Sur, Cebu, Dinagat, Leyte, Mindanao, Negros, Romblon Island Group, Samar, Sulu Archipelago (Endemic).

Figures 14 and 35.

Limnonectes macrocephalus* (Inger 1954)Rana macrodon* Boulenger 1882 (partim)*Rana magna* Stejneger 1909 (partim)*Rana macrodon macrocephala* Inger 1954*Rana magna macrocephala*, Inger 1958*Rana* (*Euphlyctis*) *magna macrocephala*, Dubois 1981*Euphlyctis magna macrocephala*, Poynton and Broadley 1985*Limnonectes* (*Limnonectes*) *macrocephalus*, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Luzon, Tayabas Province, Sampaloc (FMNH 40519).

Philippine distribution: Alabat, Cagraray, Catanduanes, Luzon, Marinduque, Masbate, Polillo, Rapu-Rapu (Endemic).

Figures 14 and 35.

Limnonectes magnus* (Stejneger 1910)Rana macrodon* Boulenger 1882 (partim)*Rana magna* Stejneger 1910; Boulenger 1920*Rana modesta* Roux 1918*Rana macrodon blythii* Boulenger, 1920 (partim)*Rana modesta magna*, Smith 1927*Rana macrodon magna*, Inger 1954 (partim)*Rana magna magna*, Inger 1958 (partim)*Rana* (*Euphlyctis*) *magna*, Dubois 1981 (partim)*Euphlyctis magna*, Poynton and Broadley 1985 (partim)*Limnonectes* (*Limnonectes*) *magnus*, Dubois 1987 “1986” (partim)

Type locality and holotype specimen: Philippines, Mindanao, Mount Apo, between Todaya and camp (4,000–6,000 ft. elev.) (USNM 35231).

Philippine distribution: Basilan, Biliran, Bohol, Camiguin Sur, Dinagat, Leyte, Mindanao, Samar (Endemic).

Figures 14 and 36.

***Limnonectes micrixalus* (Taylor 1923)**

Rana micrixalus Taylor 1923; Inger 1954; Dubois 1987 “1986”

Limnonectes micrixalus, Slevin and Leviton 1956

Type locality and holotype specimen: Philippines, Basilan, Abungabung (CAS 60143).

Philippine distribution: Basilan, Mindanao (Endemic).

Figure 15.

***Limnonectes palavanensis* (Boulenger 1894)**

Rana palavanensis Boulenger 1894; Boulenger 1920; Inger and Voris 1988

Rana microdisca palavanensis, Inger 1954

Rana (Euphlyctis) microdisca palavanensis, Dubois 1981

Euphlyctis palavanensis, Poynton and Broadley 1985

Limnonectes (Limnonectes) palavanensis, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Palawan Island (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Palawan (Non-endemic).

Figures 15 and 36.

***Limnonectes parvus* (Taylor 1920)**

Rana parva Taylor 1920; Inger 1954

Rana microdisca parva Inger 1966

Rana (Euphlyctis) microdisca parva Dubois 1981

Euphlyctis microdisca parva Poynton and Broadley 1985

Limnonectes (Limnonectes) parvus, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Mindanao, Agusan Province, Bunawan (CM 3241).

Philippine distribution: Basilan, Mindanao (Endemic).

Figures 15 and 36.

***Limnonectes visayanus* (Inger 1954)**

Rana macrodon Boulenger 1882

Rana macrodon blythii (part) Boulenger 1920

Rana magna Taylor 1923; Brown and Alcala 1970

Rana macrodon visayanus, Inger 1954

Rana magna visayanus, Inger 1958

Limnonectes (Limnonectes) visayanus, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Siquijor Island (FMNH 61636).

Philippine distribution: Bohol, Calagna-an, Cebu, Guimaras, Masbate, Negros, Panay, Poro, Romblon Island Group, Sicogon, Siquijor, Ticao (Endemic).

Figures 15 and 36.

***Limnonectes woodworthi* (Taylor 1923)**

Rana woodworthi Taylor 1923

Limnonectes (Limnonectes) woodworthi, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Luzon, Laguna Province, near Los Baños (CAS 61000).

Philippine distribution: Camiguin Norte, Catanduanes, Luzon, Polillo (Endemic).
Figures 16 and 36.

***Occidozyga diminutiva* (Taylor 1922)**

Micrixalus diminutiva Taylor 1922; Alcalá and Brown 1998

Staurois diminutives, Forcart 1946

Ooeidozyga diminutives, Inger 1954

Occidozyga diminutiva, Dubois 1981

Phrynoglossus diminutives, Dubois 1987 “1986”

Phrynoglossus diminutiva, Inger 1999; Fei, Ye, and Jiang 2010

Type locality and holotype specimen: Philippines, Mindanao, Zamboanga, “near Pasananka” (CAS 61842).

Philippine distribution: Basilan, Mindanao, Sulu Archipelago (Endemic).
Figures 16 and 36.

***Occidozyga laevis* (Günther 1858)**

Oxyglossus laevis Günther 1858; Bourret 1927

Phrynoglossus laevis, Peters 1867

Oxyglossis laevis, Smith 1916

Oxydozyga laevis, Mertens 1927

Ooeidozyga laevis, Smith 1927

Oxydozyga laevis laevis, Mertens 1930

Phrynoglossus laevis laevis, Mertens 1934

Ooeidozyga laevis laevis, Inger 1954

Occidozyga laevis, Dubois 1981; Alcalá and Brown 1998

Type locality and holotype specimen: Philippines (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Alabat, Balabac, Bohol, Bonoon, Busuanga, Cagraray, Calagna-an, Calauit, Camiguin Sur, Catanduanes, Cebu, Coron, Dinagat, Guimaras, Inampulugan, Leyte, Lubang, Luzon, Marinduque, Masbate, Mindanao, Mindoro, Negros, Palawan, Panay, Polillo, Romblon Island Group, Samar, Sicogon, Sulu Archipelago (Non-endemic).

Figures 16 and 36.

Family Eleutherodactylidae

***Eleutherodactylus planirostris* (Cope, 1862)**

Hylodes planirostris Cope 1862

Eleutherodactylus planirostris, Stejneger 1904

Eleutherodactylus ricordii planirostris, Shreve 1945

Eleutherodactylus planirostris planirostris, Schwartz 1965

Eleutherodactylus (Euhyas) planirostris, Hedges 1989; Heinicke, Duellman, and Hedges 2007

Euhyas planirostris, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, Green, and Wheeler 2006

Type locality and holotype specimen: New Providence Island, Bahamas (25.024936°N, 77.467209°W) (Peabody Essex Museum, presumed lost).

Philippine distribution: Luzon, Mindanao (Introduced; Olson et al. 2014; Sy et al. 2015).
Figure 16.

Family Megophryidae

***Leptobrachium lumadorum* Brown, Siler, Diesmos, and Alcala 2009**

Leptobrachium lumadorum Brown, Siler, Diesmos, and Alcala 2010 “2009”

Type locality and holotype specimen: Philippines, Mindanao, Zamboanga Del Sur Province, Zamboanga City, Barangay Baluno, Pasonanca Natural Park, Sitio km 24 (7.108°N, 122.0289°E; WGS84) (PNM 9561).

Philippine distribution: Basilan, Dinagat, Mindanao (Endemic).

Figures 17 and 36.

***Leptobrachium mangyanorum* Brown, Siler, Diesmos, and Alcala 2009**

Leptobrachium mangyanorum Brown, Siler, Diesmos, and Alcala 2010 “2009”

Type locality and holotype specimen: Philippines, Mindoro, Mindoro Oriental Province, Municipality of Victoria, Barangay Loyal, Sitio Panguisan, Panguisan River (13.150104°N, 121.200246°E; WGS84) (PNM 9559).

Philippine distribution: Mindoro, Semirara (Endemic).

Figures 17 and 37.

***Leptobrachium tagbanorum* Brown, Siler, Diesmos, and Alcala 2009**

Leptobrachium tagbanorum Brown, Siler, Diesmos, and Alcala 2010 “2009”

Type locality and holotype specimen: Philippines, Palawan, Palawan Province, Municipality of Puerto Princesa City, Barangay Irawan, Irawan Watershed (9.8333°N, 118.650°E; WGS84) (PNM 9560).

Philippine distribution: Palawan (Endemic).

Figures 17 and 37.

***Megophrys ligayae* Taylor 1920**

Megalophrys ligayae Taylor 1920; Inger 1999

Megophrys monticola ligayae, Inger 1954

Megophrys ligayae, Iskandar 1998

Type locality and holotype specimen: Philippines, northern Palawan (CM 3304, now CM 84521).

Philippine distribution: Balabac, Palawan (Endemic).

Figures 17 and 37.

***Megophrys stejnegeri* Taylor 1920**

Megophrys stejnegeri Taylor 1920

Megophrys monticola stejnegeri, Inger 1954

Megophrys stejnegeri, Iskandar 1998

Type locality and holotype specimen: Philippines, Mindanao, Agusan Province, Bunawan (CM 3394).

Philippine distribution: Bohol, Dinagat, Leyte, Mindanao, Samar (Endemic).

Figures 18 and 37.

Family Microhylidae***Chaperina fusca* Mocquard 1892***Chaperina fusca* Mocquard 1892; Inger, 1954*Microhyla leucostigma* Boulenger 1899*Chaperina beyeri* Taylor 1920*Nectophryne picturata* Smith 1921*Sphenophryne fusca*, Van Kampen 1923; Nieden, 1926*Sphenophryne beyeri* Van Kampen 1923*Sphenophryne leucostigma* Smith 1925

Type locality and holotype specimen: Borneo, Sintang (MNHNP 91-49).

Philippine distribution: Basilan, Mindanao, Palawan, Sulu Archipelago (Non-endemic).

Figures 18 and 37.

Kalophrynus sinensis* Peters 1867Calophrynus pleurostigma* var. *sinensis* Peters 1867*Kalophrynus sinensis*, Zug 2015

Type locality and holotype specimen: “Hongkong” [in error] (ZMB 5696). (NB: See Frost, 2015, for additional details of the complex synonymy associated with this nominal species.)

Philippine distribution: Basilan, Bohol, Camiguin Sur, Culion, Dinagat, Leyte, Mindanao, Samar (Non-endemic).

Figures 18 and 37.

Kaloula baleata* (Müller in Van Oort and Müller 1836)Bombinator baleatus* Müller in Van Oort and Müller 1836*Hyladactylus baleatus*, Tschudi 1838*Hylaedactylus baleatus*, Duméril and Bibron 1841*Hylaedactylus balteatus*, Lichtenstein and Martens 1856*Hylaedactylus baleatus* var. *concatenata* Lichtenstein and Martens 1856*Hylaedactylus lividus* Bleeker 1857*Bombinator* (*Hylaedactylus*) *baleatus*, Schlegel 1858*Hylaedactylus celebensis* Günther 1859 “1858”*Kaloula baleata*, Günther 1859 “1858”; Barbour 1909*Callula baleata*, Cope 1867; Boulenger 1882*Calohyla celebensis*, Peters 1872*Plectropus baleatus*, Knauer 1883*Kaloula baleata baleata*, Inger 1954*Kaloula baleata ghoshi*, Cherchi 1954

Type locality and holotype specimen: Indonesia, Java, Krawang (RMNH 22118).

Philippine distribution: Palawan (Non-endemic).

Figure 18.

Kaloula conjuncta* (Peters 1863)Hylaedactylus* (*Holonectes*) *conjunctus* Peters 1863*Callula conjuncta*, Cope 1867*Kaloula conjuncta*, Taylor 1920*Kaloula negrosensis* Taylor 1922*Kaloula conjuncta conjuncta*, Inger 1954 (partim)

Kalaoula conjuncta negrosensis, Inger 1954 (partim)

Kaloula conjuncta stickeli, Inger 1954

Kaloula conjuncta meridionalis, Inger 1954 (partim)

Type locality and holotype specimen: Philippines, Luzon Island (NMW 22888).

Philippine distribution: Alabat, Borocay, Caluya, Catanduanes, Cebu, Guimaras, Leyte, Luzon, Mindanao, Mindoro, Negros, Pacijan, Panay, Polillo, Poro, Romblon Island Group, Semirara, Siquijor, Sulu Archipelago (Endemic).

Figures 19 and 37.

***Kaloula kalingensis* Taylor, 1922**

Kaloula kalingensis Taylor 1922; Ross and Gonzales 1992

Kaloula baleata kalingensis, Inger 1954

Type locality and holotype specimen: Philippines, Luzon, Mountain Province, Kalinga, Balbalan (CAS 61462).

Philippine distribution: Luzon, Palau (Endemic).

Figures 19 and 37.

***Kaloula kokacii* Ross and Gonzales 1992**

Kaloula kokacii Ross and Gonzales 1992

Type locality and holotype specimen: Philippines, Catanduanes, Gigmoto Municipality, from abaca near the Buadan River, Summit Boradan (8.5 km W and 1 km N of Gigmoto) (13.8°N, 124.316667°E; WGS84; 200–300 m elev.) (PNM 2043).

Philippine distribution: Catanduanes, Luzon (Endemic).

Figure 19.

***Kaloula picta* (Duméril and Bibron 1841)**

Plectropus pictus Duméril and Bibron 1841; Guibé 1950

Kaloula picta, Günther 1859; Parker 1934

Callula picta, Günther 1864

Type locality and holotype specimen: Philippines, Luzon, Manille (= Manila) (MNHNP 5027).

Philippine distribution: Alabat, Babuyan Island Group, Bohol, Caluya, Camiguin Sur, Catanduanes, Cebu, Cocomo, Cuyo, Dinagat, Guimaras, Lapinig Grande, Leyte, Lubang, Luzon, Mactan, Mindanao, Mindoro, Negros, Palawan, Panay, Polillo, Ponson, Rapu-Rapu, Samar, Semirara, Verde (Endemic).

Figures 19 and 38.

***Kaloula pulchra* Gray 1831**

Kaloula pulchra Gray 1831; Barbour 1909

Hylaedactylus bivittatus Cantor 1847; Boulenger 1882; Günther, 1859 “1858”; Bourret 1942

Callula pulchra, Günther 1864

Caloula pulchra, Stoliczka 1870

Calohyla pulchra, Peters and Doria 1878

Callula macrodactyla Boulenger, 1887; Parker 1934; Bourret 1942

Callula (Kallula) pulchra, Bourret 1927

Kaloula pulchra pulchra, Parker 1934

Kaloula pulchra hainana, Gressitt 1938

Kaloula pulchra macrocephala, Bourret 1942

Kaloula macrocephala Ohler 2003

Type locality and holotype specimen: China (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Cebu, Luzon, Mindoro, Palawan (Introduced; Diesmos et al. 2006; Sy et al. 2014).

Figures 20 and 38.

***Kaloula rigida* Taylor 1922**

Kaloula rigida Taylor 1922; Parker 1934; Slevin and Leviton 1956

Type locality and holotype specimen: Philippines, Luzon, Mountain Province, Kalinga, Balbalan (CAS 61475).

Philippine distribution: Luzon (Endemic).

Figures 20 and 38.

***Kaloula walteri* Diesmos, Brown, and Alcala 2002**

Kaloula walteri Diesmos, Brown, and Alcala 2002

Type locality and holotype specimen: Philippines, Luzon, Quezon Province, Barangay Lalo, Municipality of Tayabas, on the southeast slope of Mt. Banahao (14.066667°N, 121.483333°E; WGS84; 950 m elev.) (PNM 6725).

Philippine distribution: Luzon, Polillo (Endemic).

Figures 20 and 38.

***Microhyla petrigena* Inger and Frogner 1979**

Microhyla (Microhyla) petrigena, Dubois 1987

Type locality and holotype specimen: Malaysia, Borneo, Sarawak, Kapit District, Nanga Tekalit. (FMNH 207705).

Philippine distribution: Tawi-Tawi (Non-endemic).

Figures 20 and 38.

***Oreophryne anulata* (Stejneger 1908)**

Phrynixalus anulatus Stejneger 1908; Taylor 1920

Chaperina visaya Taylor 1920

Phrynixalus annulatus Taylor 1920

Oreophryne annulata, Parker 1934; Inger 1954

Type locality and holotype specimen: Philippines, Mindanao, Davao (USNM 35399).

Philippine distribution: Leyte, Mindanao, Samar (Endemic).

Figures 21 and 38.

***Oreophryne nana* Brown and Alcala 1967**

Oreophryne nana Brown and Alcala 1967

Type locality and holotype specimen: Philippines, Camiguin, Mt. Hibok-hibok, on the northwest side of Nacawa volcano (1,800–3,000 ft. elev.) (CAS-SU 22055).

Philippine distribution: Camiguin Sur (Endemic).

Figures 21 and 38.

Family Ranidae

Amnirana nicobariensis (Stoliczka 1870)

Hyla bilineata Van-Ernest *in* Daudin 1800; Daudin *in* Sonnini de Manoncourt and Latreille 1801

Calamita bilineatus Merrem 1820

Auletris bilineatus, Wagler 1830

Hylorana nicobariensis Stoliczka 1870; Deckert 1938

Rana macularia var. *javanica* Horst 1883

Rana javanica, Boulenger 1884; Van Kampen 1907

Rana nicobariensis, Boulenger 1885

Rana erythraea var. *elongate*, Werner 1892

Rana lemniscata Boettger 1893

Rana (*Hylorana*) *nicobariensis*, Boulenger 1920

Rana sanchezi Taylor 1920

Rana suluensis Taylor 1920

Rana (*Hylorana*) *nicobariensis*, Boulenger 1920; Van Kampen 1923

Rana nicobariensis javanica, Mertens 1927

Rana nicobariensis nicobariensis, Inger 1954

Rana (*Sylvirana*) *nicobariensis*, Dubois 1992

Rana nicobariensis, Alcala and Brown 1998

Sylvirana nicobariensis, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana nicobariensis, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Amnirana nicobariensis, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Indonesia, Java (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Palawan, Sulu Archipelago (Non-endemic).

Figure 21.

Hylarana erythraea (Schlegel 1837)

Hyla erythraea Schlegel 1837

Hylarana erythraea, Tschudi 1838; Bourret 1942; Fei, Ye, and Huang 1990; Chen, Murphy, Lathrop, Ngo, Orlov, Ho, and Somorjai 2005

Limnodytes erythraeus, Duméril and Bibron 1841

Hylorana erythraea, Günther 1864; Deckert 1938

Rana erythraea, Boulenger 1882

Rana (*Hylorana*) *erythraea*, Boulenger *in* Mason 1882

Rana (*Hylarana*) *erythraea*, Müller 1887

Rana (*Limnodytes*) *erythraea*, Bourret 1927

Type locality and holotype specimen: Sumatra (RMNH 1744, 1746, 1749; MNHNP 4570–4572, syntypes).

Philippine distribution: Borocay, Calagna-an, Guimaras, Leyte, Luzon, Masbate, Mindoro, Negros, Panay, Romblon Island Group, Samar (Introduced; Diesmos et al. 2006).

Figures 21 and 38.

Lithobates catesbeianus* (Shaw 1802)Rana catesbeiana* Shaw, 1802; Boulenger 1920; Dubois 1987 “1986”*Rana pipiens* Daudin 1802*Rana taurina* Cuvier 1817*Rana mugiens* Merrem, 1820*Rana scapularis* Harlan, 1826*Rana conspersa* Le Conte 1855*Rana catesbyana* Cope 1889; Werner 1909*Rana (Rana) catesbeiana*, Boulenger, 1920*Rana nantaiwuensis* Hsü 1930*Rana mugicus* Angel 1947*Rana (Aquarana) catesbeiana*, Dubois, 1992; Hillis 2007*Rana (Novirana, Aquarana) catesbeiana*, Hillis and Wilcox 2005*Lithobates catesbeianus*, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, Green, and Wheeler, 2006*Lithobates (Aquarana) catesbeianus*, Dubois 2006

Type locality and holotype specimen: North America (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Leyte, Luzon, Mindanao, Mindoro, Panay (Introduced; Diesmos et al. 2006).

Figures 22 and 39.

Pulchrana grandocula* (Taylor 1920)Rana grandocula* Taylor 1920; Inger and Tan 1996*Rana philippinensis* Taylor 1920*Rana yakani* Taylor 1922*Rana signata gradocula*, Inger 1954; Alcalá and Brown 1998*Rana signata*, Frost 1985*Rana (Pulchrana) grandocula*, Dubois 1992*Pulchrana grandocula*, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006*Hylarana grandocula*, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007; Brown and Siler 2013*Pulchrana grandocula*, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Philippines, Mindanao, Agusan Province, near Bunawan (CM 3501).

Philippine distribution: Basilan, Biliran, Bohol, Camiguin Sur, Dinagat, Leyte, Mindanao, Samar (Endemic).

Figures 22 and 39.

Pulchrana guttmani* (Brown 2015)Rana grandocula* Brown and Guttman 2002 (partim)*Hylarana* sp. 2, Brown and Siler 2013*Hylarana guttmani* Brown 2015

Type locality and holotype specimen: Philippines, southern Mindanao Island, South Cotabato Province (~2 km north of border with Sarangani Province), Municipality of Kiamba,

Barangay Badtasan, Sitio Banate, Mt. Busa (6.0923°N, 124.6709°E; WGS84; 1,200 m elev.) (PNM 9790, formerly KU 326399).

Philippine distribution: Mindanao (Endemic).

Figure 43.

***Pulchrana mangyanum* (Brown and Guttman 2002)**

Rana mangyanum Brown and Guttman 2002

Pulchrana mangyanum, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana mangyanum, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007; Brown and Siler 2013

Pulchrana mangyanum, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Philippines, Mindoro, Oriental Mindoro Province, Municipality of Puerto Galera (within 1 km of the border of the Municipality of San Teodoro), Barangay Villaflor (15 km from Puerto Gallera City on Puerto Gallera-Calapan Road), Tamaraw Falls (unnamed river) (150 m elev.) (PNM 6270).

Philippine distribution: Mindoro, Semirara (Endemic).

Figures 22 and 39.

***Pulchrana melanomenta* (Taylor 1920)**

Rana melanomenta Taylor 1920; Brown and Gittman 2002

Rana (Pulchrana) melanomenta, Dubois 1992

Pulchrana melanomenta, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana melanomenta, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007; Brown and Siler 2013

Pulchrana melanomenta, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Philippines, Sulu Archipelago, Papahag Island (Bur. Sci. Manila No. 1661, destroyed).

Philippine distribution: Papahag (Endemic).

Figure 22.

***Pulchrana moellendorffi* (Boettger 1893)**

Rana moellendorffi Boettger 1893; Brown and Gittman 2002

Rana (Hylarana) moellendorffi, Boulenger 1920

Rana signata moellendorffi, Inger 1954; Alcala and Brown 1998

Rana (Hylarana) moellendorffi, Dubois 1987 “1986”

Rana (Pulchrana) moellendorffi, Dubois 1992

Pulchrana moellendorffi, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana moellendorffi, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007; Brown and Siler 2013

Pulchrana moellendorffi, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Philippines, Calamianes Island Group, Culion Island (SMF 5432).

Philippine distribution: Balabac, Busuanga, Caluit, Coron, Culion, Palawan (Endemic).
Figures 23 and 39.

***Pulchrana similis* (Günther 1873)**

Polypedates similis Günther 1873

Rana similis, Boulenger 1882; Brown and Guttman 2002

Rana (Hylarana) signata Boulenger 1920

Rana signata similis, Inger 1954; Alcala and Brown 1998

Rana (Pulchrana) similis, Dubois 1992

Pulchrana similis, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana similis, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007; Brown and Siler 2013

Pulchrana similis, Oliver, Prendini, Kraus, and Raxworthy 2015

Type locality and holotype specimen: Philippines, Luzon, Laguna del Bay (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Cagraray, Catanduanes, Luzon, Polillo, Rapu-Rapu (Endemic).
Figures 23 and 39.

***Sanguirana albotuberculata* (Inger 1954)**

Rana everetti albotuberculata Inger 1954

Rana (Chalcorana) albotuberculata, Dubois 1992

Rana albotuberculata, Brown, McGuire, and Diesmos 2000

Hydrophylax albotuberculata, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana albotuberculata, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Chalcorana albotuberculata, Fei, Ye, and Jiang 2010

Sanguirana albotuberculata, Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011

Type locality and holotype specimen: Philippines, Leyte, Cabalian (MCZ 23190).

Philippine distribution: Leyte, Mindanao, Samar (Endemic).
Figures 23 and 39.

***Sanguirana aurantipunctata* Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011**

Type locality and holotype specimen: Philippines, Luzon, Nueva Vizcaya Province, Municipality of Quezon, Barangay Maddiangat, Sitio Parola, Mt. Palali (16.438°N, 121.225°E; WGS84; 1,500 m elev.) (PNM 9727).

Philippine distribution: Luzon (Endemic).
Figures 23 and 39.

***Sanguirana everetti* (Boulenger 1882)**

Rana everetti Boulenger 1882; Brown, McGuire, and Diesmos 2000

Rana mearnsi Stejneger 1905

Rana dubita Taylor 1920

Rana (Hylarana) everetti, Boulenger 1920

Rana (Hylarana) mearnsi Boulenger 1920

Rana merrilli Taylor 1922

Rana (Hylarana) everetti, Van Kampen 1923; Dubois 1987 “1986”

Rana everetti everetti, Inger 1954

Rana (Chalcorana) everetti, Dubois 1992

Hydrophylax everetti, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana everetti, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Chalcorana everetti, Fei, Ye, and Jiang 2010

Sanguirana everetti, Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011

Type locality and holotype specimen: Philippines, Mindanao, Zamboanga (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Mindanao (Endemic).

Figures 24 and 39.

***Sanguirana igorota* (Taylor 1922)**

Rana igorota Taylor 1922; Brown, McGuire, and Diesmos 2000

Rana everetti luzonensis Inger 1954

Rana (Chalcorana) luzonensis, Dubois 1992

Hydrophylax igorata, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana igorota, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Chalcorana igorota, Fei, Ye, and Jiang 2010

Sanguirana igorata, Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011

Type locality and holotype specimen: Philippines, Luzon, Kalinga Subprovince, Balbalan (CAS 61484).

Philippine distribution: Luzon (Endemic).

Figures 24 and 40.

***Sanguirana luzonensis* (Boulenger 1896)**

Rana luzonensis Boulenger 1896; Dubois 1987 “1986”; Brown, McGuire, and Diesmos 2000

Rana (Hylarana) luzonensis, Boulenger 1920

Rana guerreroi Taylor 1920

Rana merilli Taylor 1922

Rana igorata Taylor 1922

Rana tafti Taylor 1922

Rana everetti luzonensis, Inger 1954; Alcala and Brown 1998

Rana (Chalcorana) luzonensis, Dubois 1992

Hydrophylax luzonensis, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana luzonensis, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Chalcorana luzonensis, Fei, Ye, and Jiang 2010

Sanguirana luzonensis, Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011

Type locality and holotype specimen: Philippines, N. Luzon Highlands of Lepauto (= Lepanto) (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Luzon, Catanduanes, Marinduque, Polillo (Endemic).

Figures 24 and 40.

***Sanguirana sanguinea* (Boettger 1893)**

Rana sanguinea, Boettger 1893; Inger 1954

Rana varians Boulenger 1894

Rana (Hylarana) sanguinea, Boulenger 1920

Rana (Hylarana) varians, Boulenger 1920

Hylarana varians, Deckert 1938

Rana (Hylarana) sanguinea, Dubois 1987 “1986”

Rana (Hylarana) varians, Dubois 1987 “1986”

Hylarana (Hylarana) varians, Fei, Ye, and Huang 1990

Rana (Sanguirana) sanguinea, Dubois 1992

Rana (Sanguirana) varians, Dubois 1992

Hylarana sanguinea, Song, Jang, Zou, and Shi 2002

Sanguirana sanguinea, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Sanguirana varians Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Type locality and holotype specimen: Philippines, Calamianes Island Group, Culion Island (SMF 1062a, now SMF 6221).

Philippine distribution: Busuanga, Culion, Palawan (Endemic).

Figures 24 and 40.

***Sanguirana tipanan* (Brown, McGuire, and Diesmos 2000)**

Rana tipanan Brown, McGuire, and Diesmos 2000

Hydrophylax tipanan, Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, and Green 2006

Hylarana tipanan, Che, Pang, Zhao, Wu, Zhao, and Zhang 2007

Chalcorana tipanan, Fei, Ye, and Jiang 2010

Sanguirana tipanan, Fuiten, Welton, Diesmos, Barley, Oberheide, Duya, Rico, and Brown 2011

Type locality and holotype specimen: Philippines, Luzon, Aurora Province, Municipality of San Luis, Aurora National Park, 1.2 km S, 1.3 km E of Barangay Villa Aurora, east side of Mt. Ma-aling-aling in the Kabatangan river drainage (15.651667°N, 121.351944°E; 470 m elev.) (PNM 5727).

Philippine distribution: Luzon (Endemic).

Figures 25 and 40.

***Staurois natator* (Günther 1858)**

Ixalus natator Günther 1858

Ixalus guttatus Günther 1858

Staurois natator, Cope 1865; Inger 1954

Rana natatrix Boulenger 1882

Ixalus granulatus Boettger 1888

Staurois nubilus, Boulenger 1918

Rhacophorus granulatus Ahl 1927

Rana guttatus Smith 1931

Type locality and holotype specimen: Philippines (BMNH 1933.9.19.10.9–11, syntypes).

Philippine distribution: Basilan, Biliran, Bohol, Dinagat, Leyte, Mindanao, Samar (Endemic).

Figures 25 and 40.

***Staurois nubilus* (Mocquard 1890)**

Ixalus nubilus Mocquard 1890

Ixalus natator var. *nubilus*, Mocquard 1892; Guibé 1950

Staurois nubilus, Boulenger 1918; Decker 1938; Inger and Tan 1996

Type locality and holotype specimen: Philippines, Palawan (MNHNP 1889.344–46, syntypes).

Philippine distribution: Busuanga, Culion, Palawan (Endemic).

Figures 25 and 40.

Family Rhacophoridae

***Kurixalus appendiculatus* (Günther 1858)**

Polypedates appendiculatus Günther 1858

Rhacophorus appendiculatus, Boulenger 1882; Ahl 1931; Brown and Alcalá 1994; Harvey, Pemberton, and Smith 2002

Rhacophorus phyllopygus Werner 1900

Rhacophorus chaseni Smith 1924

Rhacophorus appendiculatus chaseni, Smith 1930

Rhacophorus appendiculatus appendiculatus, Smith 1930; Inger 1954

Rhacophorus (Rhacophorus) chaseni, Ahl 1931

Rhacophorus (Rhacophorus) appendiculatus appendiculatus, Wolf 1936

Leptomantis appendiculatus, Iskandar and Colijn 2000

Kurixalus appendiculatus, Yu, Zhang, and Yang 2013

Type locality and holotype specimen: Philippines (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Basilan, Bohol, Camiguin Sur, Leyte, Luzon, Mindanao, Samar (Non-endemic).

Figures 25 and 40.

***Nyctixalus pictus* (Peters 1871)**

Ixalus pictus Peters 1871

Rhacophorus anodon Van Kampen 1907

Philautus pictus, Barbour 1912

Philautus anodon, Van Kampen 1923

Rhacophorus (Philautus) anodon, Ahl 1931

Rhacophorus (Philautus) pictus, Ahl 1931

Hazelia picta, Taylor 1962

Philautus pictus pictus, Inger 1966

Hazelia anodon, Liem 1970

Nyctixalus anodon, Dubois 1981

Edwardtayloria picta, Dring 1982

Edwardtayloria picta, Alcala 1986

Nyctixalus pictus, Brown and Alcala 1994; Matsui 1996

Type locality and holotype specimen: Malaysia (Borneo), Sarawak (MSNG 10062).

Philippine distribution: Palawan (Endemic).

Figure 26.

***Nyctixalus spinosus* (Taylor 1920)**

Hazelia spinosa Taylor 1920

Rhacophorus (Philautus) spinosus, Ahl 1931

Rhacophorus leprosus spinosus, Wolf 1936

Philautus spinosus, Inger 1954

Hazelia spinosa, Liem 1970

Edwardtayloria spinosa, Marx 1975

Nyctixalus spinosus, Dubois 1981; Brown and Alcala 1994

Edwardtayloria spinosa, Alcala 1986

Type locality and holotype specimen: Philippines, Mindanao, Agusan Province, Bunawan (CM 3420).

Philippine distribution: Basilan, Bohol, Leyte, Mindanao, Samar (Endemic).

Figures 26, 40, and 41.

***Philautus acutirostris* (Peters 1867)**

Ixalus acutirostris Peters 1867

Philautus acutirostris, Stejneger 1905; Inger 1954

Philautus woodi Stejneger 1905

Philautus basilanensis Taylor 1922

Rhacophorus (Philautus) woodi Ahl 1931

Rhacophorus (Philautus) basilanensis Ahl 1931

Rhacophorus (Philautus) acutirostris, Ahl 1931

Philautus (Philautus) acutirostris, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, eastern Mindanao (NMW 22885, ZMB 5690, syntypes).

Philippine distribution: Basilan, Bohol, Mindanao (Endemic).

Figures 26 and 41.

***Philautus everetti* (Boulenger 1894)**

Rhacophorus everetti Boulenger 1894

Polypedates everetti, Taylor 1920

Rhacophorus (Rhacophorus) buergeri everetti, Wolf 1936

Rhacophorus everetti everetti, Inger 1954

Philautus everetti, Hertwig, Das, Schweizer, Brown, and Haas 2012

Type locality and holotype specimen: Philippines, Palawan (BMNH 94.6.3.126–127, syntypes).

Philippine distribution: Palawan (Endemic).

Figures 26 and 41.

***Philautus leitensis* (Boulenger 1897)**

Ixalus leitensis Boulenger 1897

Philautus leitensis, Stejneger 1905; Inger 1954; Bossuyt and Dubois 2001

Rhacophorus (Philautus) leitensis, Ahl 1931

Type locality and holotype specimen: Philippines, Leyte (BMNH 96.12.11.92).

Philippine distribution: Bohol, Leyte, Mindanao, Samar (Endemic).

Figures 27 and 41.

***Philautus longicrus* (Boulenger 1894)**

Ixalus longicrus Boulenger 1894

Philautus longicrus, Stejneger 1905; Inger 1954

Rhacophorus (Philautus) longicrus, Ahl 1931

Philautus (Philautus) longicrus, Bossuyt and Dubois 2001

Type locality and holotype specimen: Philippines, Palawan (BMMH 94.6.30.129–131, now BMMH 1947.2.6.28–30, syntypes).

Philippine distribution: Palawan (Non-endemic).

Figures 27 and 41.

***Philautus poecilus* Brown and Alcala 1994**

Philautus poecilus Brown and Alcala 1994

Philautus (Philautus) poecilus, Bossuyt and Dubois 2001

Type locality and holotype specimen: Philippines, Mindanao, Agusan del Norte Province, south side of Mt. Hilong-hilong (CAS 133526).

Philippine distribution: Mindanao (Endemic).

Figures 27 and 41.

***Philautus schmackeri* (Boettger 1892)**

Ixalus schmackeri Boettger 1892

Ixalus mindorensis Boulenger 1897

Philautus schmackeri, Stejneger 1905; Taylor 1920; Inger 1954; Dubois 1987 “1986”

Philautus mindorensis Taylor 1920; Stejneger 1905

Rhacophorus (Philautus) schmackeri, Ahl 1931

Rhacophorus (Philautus) mindorensis Ahl 1931

Type locality and holotype specimen: Philippines, Mindoro, Mt. Halcone (SMF 1099a, now SMF 7035).

Philippine distribution: Mindoro (Endemic).

Figure 27.

***Philautus surdus* (Peters 1863)**

Polypedates surdus Peters 1863

Rhacophorus surdus, Boulenger 1882

Philautus williamsi Taylor 1922

Rhacophorus (Philautus) williamsi, Ahl 1931

Rhacophorus (Rhacophorus) surdus, Ahl 1931

Rhacophorus (Rhacophorus) buergeri surdus, Wolf 1936

Rhacophorus lissobrachius Inger 1954

Rhacophorus surdus, Inger 1954

Philautus surdus, Liem 1970; Brown and Alcala 1994

Philautus lissobrachius, Liem 1970

Philautus (Philautus) lissobrachius, Dubois 1987 “1986”

Philautus (Philautus) surdus, Dubois 1987 “1986”

Type locality and holotype specimen: Philippines, Luzon (ZMB 4920).

Philippine distribution: Bohol, Luzon, Mindanao (Endemic).

Figures 28 and 41.

***Philautus surrufus* Brown and Alcalá 1994**

Rhacophorus surdus Rabor and Alcalá 1959 (partim)

Philautus surrufus, Brown and Alcalá 1994

Philautus (Philautus) surrufus, Bossuyt and Dubois 2001

Type locality and holotype specimen: Philippines, Mindanao, Misamis Occidental Province, about 10 km SE of Masawan, on the west side of Dapitan Peak (1,800–1,900 m elev.) (CAS-SU 21013).

Philippine distribution: Mindanao (Endemic).

Figure 28.

***Philautus worcesteri* (Stejneger 1905)**

Cornufer worcesteri Stejneger 1905

Rhacophorus emembranatus Inger 1954

Philautus emembranatus Liem 1970

Philautus (Philautus) emembranatus, Dubois 1987 “1986”

Philautus worcesteri, Brown, Alcalá, and Brown 1998

Type locality and holotype specimen: Philippines, Mindanao, Mount Apo (6,000 ft. elev.) (USNM 34784).

Philippine distribution: Mindanao (Endemic).

Figures 28 and 41.

***Polypedates leucomystax* (Gravenhorst 1829)**

Hyla leucomystax Gravenhorst 1829

Hyla sexvirgata Gravenhorst 1829

Hyla quadrilineata Wiegmann 1834

Polypedates leucomystax, von Tschudi 1838

Hyla leucopogon von Tschudi 1838

Hyla quadrivirgata von Tschudi 1838

Polypedates rugosus Duméril and Bibron 1841

Polypedates quadrilineatus, Günther 1859 “1858”

Limnodytes celebensis Fitzinger 1861 “1860”

Polypedates hecticus Peters 1863; Taylor 1920

Rhacophorus hecticus, Boulenger 1882

Rhacophorus maculatus var. *quadrilineata*, Boulenger 1882

Hylorana longipes Fischer 1885

Polypedates maculatus quadrilineatus, Fischer 1885

Rhacophorus maculatus Boettger 1886

Rhacophorus leucomystax, Boulenger 1889; Van Kampen 1923; Ahl 1931

Rhacophorus leucomystax leucomystax, Mocquard 1890

Rhacophorus leucomystax quadrilineatus, Mocquard 1890; Inger 1954

Rhacophorus leucomystax var. *sexvirgata*, Boettger 1894

Rhacophorus leucomystax quadrilineata, Werner 1903

Rhacophorus maculatus leucomystax, Annandale 1912
Rhacophorus maculatus himalayensis Annandale 1912
Hyla wirzi Roux 1927
Polypedates leucomystax, Taylor 1920
Rhacophorus (Polypedates) leucomystax, Bourret 1927
Rhacophorus (Polypedates) quadrilineatus, Bourret 1927
Rhacophorus kampeni Ahl 1927
Rhacophorus (Rhacophorus) hecticus Ahl 1931
Rhacophorus (Rhacophorus) himalayanus, Ahl 1931
Rhacophorus (Rhacophorus) kampeni, Ahl 1931
Rhacophorus (Rhacophorus) leucomystax leucomystax, Wolf 1936
Rhacophorus (Rhacophorus) wirzi Forcart 1946
Polypedates leucomystax, Alcala 1986
Polypedates leucomystax, Dutta 1997

Type locality and holotype specimen: Indonesia, Java (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Batan, Bohol, Cagayan, Cagraray, Calagna-an, Caluya, Camiguin Norte, Camiguin Sur, Catanduanes, Cebu, Dinagat, Fuga, Gigantes Norte, Gigantes Sur, Guimaras, Inampulugan, Jolo, Leyte, Lubang, Luzon, Mactan, Marinduque, Masbate, Mindanao, Mindoro, Negros, Pacijan, Palaui, Palawan, Pan de Azucar, Panay, Polillo, Romblon Island Group, Samar, Semirara, Sibay, Sicogan, Verde (Non-endemic).

Figures 28 and 42.

***Polypedates macrotis* (Boulenger 1891)**

Rhacophorus macrotis Boulenger 1891
Polypedates macrotis, Günther 1895; Liem 1970; Alcala 1986
Philautus montanus Taylor 1920
Polypedates linki Taylor 1922
Rhacophorus (Philautus) alticola Ahl 1931
Rhacophorus (Rhacophorus) macrotis, Ahl 1931
Rhacophorus (Rhacophorus) lincki Ahl 1931
Rhacophorus leucomystax linki, Wolf 1936; Inger 1954
Rhacophorus (Polypedates) macrotis, Bossuyt and Dubois 2001

Type locality and holotype specimen: Borneo, Sarawak, Baram district (BMNH 91.1.27.8, now BMNH 1947.2.8.18).

Philippine distribution: Busuanga, Calauit, Dumarán, Palawan, Sulu Archipelago (Non-endemic).

Figures 29 and 42.

***Rhacophorus bimaculatus* (Peters 1867)**

Leptomantis bimaculata Peters 1867
Ixalus bimaculatus, Boulenger 1882
Philautus bimaculatus, Stejneger 1905; Inger 1954
Philautus zamboangensis Taylor 1922
Rhacophorus (Philautus) bimaculatus, Ahl 1931
Rhacophorus bimaculatus, Liem 1970
Rhacophorus (Leptomantis) bimaculatus, Dubois 1987 “1986”
Leptomantis bimaculatus, Iskandar and Colijn 2000
Rhacophorus bimaculatus, Brown and Alcala 1994; Harvey, Pemberton, and Smith 2002

Type locality and holotype specimen: Philippines, Mindanao, Upper Valley of the Agusan (ZMB 5681, NHMW 16091, syntypes).

Philippine distribution: Bohol, Catanduanes, Dinagat, Leyte, Luzon, Mindanao, Polillo, Samar (Endemic).

Figures 29 and 42.

***Rhacophorus pardalis* Günther 1858**

Rhacophorus pardalis Günther 1858

Rhacophorus rizali Boettger 1897, 1899

Rhacophorus pulchellus Werner 1900

Polypedates pardalis, Taylor 1920

Rhacophorus (*Rhacophorus*) *pardalis*, Ahl 1931

Rhacophorus (*Rhacophorus*) *pulchellus*, Ahl 1931

Rhacophorus pardalis pardalis, Wolf 1936; Inger 1954

Rhacophorus pardalis pulchellus, Wolf 1936

Rhacophorus pardalis rhysocephalus, Wolf 1936

Rhacophorus rhysocephalus, Inger and Voris 2001

Type locality and holotype specimen: Philippines (Status and whereabouts of holotype unknown; not traced).

Philippine distribution: Basilan, Bohol, Camiguin Sur, Catanduanes, Dinagat, Leyte, Luzon, Mindanao, Mindoro, Negros, Romblon Island Group, Samar, Siquijor (Non-endemic).

Figures 29 and 42.

Class Amphibia, Order Gymnophiona
Family Ichthyophiidae

***Ichthyophis glandulosus* (Taylor 1923)**

Ichthyophis glandulosus Taylor 1923

Ichthyophis monochrous Inger 1954; Alcalá 1986

Type locality and holotype specimen: Philippines, Basilan, Abungabung (= Abung Abung) (CAS 60073).

Philippine distribution: Basilan, Mindanao (Endemic).

Figures 29 and 42.

***Ichthyophis mindanaoensis* (Taylor 1960)**

Ichthyophis monochrous Inger 1954; Alcalá 1986

Ichthyophis mindanaoensis Taylor 1960

Type locality and holotype specimen: Philippines, Mindanao, Davao Province, Mt. Apo, Todaya (2,800 ft. elev.) (FMNH 50958).

Philippine distribution: Mindanao (Endemic).

Figure 30.

***Ichthyophis weberi* Taylor 1920**

Ichthyophis weberi Taylor 1920

Caudacaecilia weberi, Taylor 1923, 1968

Ichthyophis monochrous Inger 1954; Alcalá 1986

Ichthyophis weberi, Nishikawa, Matsui, Yong, Ahmad, Yambun, Belabut, Sudin, Hamidy, Orlov, Ota, Yoshikawa, Tominaga, and Shimada 2012

Type locality and holotype specimen: Philippines, Palawan, Malatgan River (CAS-SU 21758, neotype).

Philippine distribution: Palawan (Endemic).

Figure 30.

CONCLUSIONS

Our understanding of biodiversity of amphibians in the Philippines has increased substantially over the last century as a result of continued faunal surveys over a greater proportion of the archipelago and, more recently, with the increased availability of genetic data guiding identification and discovery of unique evolutionary lineages (Brown et al. 2013). Vouchered global collections now exceed 43,000 specimen records, housed among more than 30 museums in seven countries. The amphibian fauna in the Philippines includes members of nine anuran families (Bombinatoridae, Bufonidae, Ceratobatrachidae, Dicroglossidae, Eleutherodactylidae, Megophryidae, Microhylidae, Ranidae, Rhacophoridae) and one gymnophionan family (Ichthyophiidae). Four of these families are represented by endemic species only in the archipelago (Bombinatoridae, Ceratobatrachidae, Ichthyophiidae, Megophryidae).

What once was considered a depauperate amphibian fauna composed of a number of widespread species distributed across larger regions of Southeast Asia (Inger 1954; Brown and Alcala 1970a, b), we take note of the fact that the diversity of endemic amphibian species in the Philippines has risen precipitously (Figure 1) in recent years. Currently, there are 112 species recorded in the archipelago, 94 of which are endemic (83.9% amphibian endemism). In contrast, truly widespread (non-endemic) species account for only 16.1% (18 species; Figure 1). Furthermore, nearly one-third of the country's non-endemic species are the result of introductions (*Eleutherodactylus planirostris*, *Hoplobatrachus rugulosus*, *Hylarana erythraea*, *Kaloula pulchra*, *Lithobates catesbeianus*, and *Rhinella marina*; Figure 1). Although the rate of discovery of non-endemic species of amphibians has slowed considerably over the last hundred years, the number of endemic species continues to grow with little indication of slowing. Within the last 20 years alone, 27 new amphibian species have been described (nearly one-quarter of the country's recognized diversity), all endemic to the Philippines (Figure 1). These data suggest that considerable cryptic diversity and underestimated regional diversity exist throughout the archipelago. Continued efforts to describe and study the archipelago's amphibian fauna are necessary for successful conservation of threatened taxa and clarification of the broader evolutionary mechanisms that drive such diversity.

ACKNOWLEDGEMENTS

We thank the many museums all over the world that shared their Philippine herpetological data with us, either directly (British Museum of Natural History, California Academy of Sciences, Natural History Museum Vienna, Naturalis Biodiversity Center, Smithsonian National Museum of Natural History, University of Texas Biodiversity Collections [formerly Texas Natural History Collections], University of Kansas Biodiversity Institute and Natural History Museum) or through web portals (GBIF, VertNet). This summary could not have been possible without open access to vouchered museum specimen data. We also thank Janalee P. Caldwell, Scott Travers, and Jason Fernandez for sharing their photos. CDS and RMB are grateful for all institutions that provided financial support for fieldwork over the last two decades, including support for ACD from the National Museum of the Philippines, and for CDS from the Sam Noble Oklahoma Museum of Natural History, a Fulbright Fellowship, a Fulbright-Hayes Fellowship, and NSF DEB 0804115 and




FIGURE 1.—Species accumulation curve for new amphibian species described in the Philippines, from 1758–2015.

IOS 1353683, and support for RMB from Miami University, University of Texas (Austin), and University of Kansas Biodiversity Institute, and NSF DEB 0743491 and EF-0334952. Members of the Siler Lab provided critical reviews of early drafts of the manuscript. We also wish to express our appreciation to our anonymous reviewers for their suggestions for the improvement of this contribution.

LITERATURE CITED

- ALCALA, A. C., W. C. BROWN, AND A. C. DIESMOS. 1998. Two new species of the genus *Platymantis* (Amphibia: Ranidae) from Luzon Island, Philippines. *Proceedings of the California Academy of Sciences*, ser. 4, 50:381–388.
- AMPHIBIAWEB. 2015. *AmphibiaWeb: Information on Amphibian Biology and Conservation*. Berkeley, CA, USA. <<http://amphibiaweb.org>>. Cited 15 October 2015.
- BARBOUR, T. 1923. The frogs of the Fiji Islands. *Proceedings of the Academy of Natural Sciences Philadelphia* 75:111–115.
- BARBOUR, T. 1938. Notes on *Nectophryne*. *Proceedings of the Biological Society of Washington* 51:191–196.
- BLEEKER, P. 1858. Bestuursvergadering, gehouden ten huize van den heer Dr Bruijn Kops den 11n Maart 1858. *Natuurkundig Tijdschrift voor Nederlandsch Indië* 16:183–192.
- BOETTGER, O. 1892. *Katalog der Batrachier-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main*. Druck von Gebrüder Knauer, Frankfurt, Germany. 73 pp.
- BOETTGER, O. 1893. *Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. I. Teil (Rhynchocephalen, Schildkröten, Krokodile, Eidechsen, Chamäleons)*. Gebrüder Knauer, Frankfurt, Germany. 140 pp.
- BOULENGER, G. A. 1882. *Catalogue of the Batrachia Salientia s. Ecaudata in the Collection of the British Museum*. Second Edition. Taylor and Francis, London, UK. 160 pp.
- BOULENGER, G. A. 1887. Descriptions of new reptiles and batrachians in the British Museum (Natural History).—Part III. *Annals and Magazine of Natural History*, ser. 5, 20:50–53.
- BOULENGER, G. A. 1891. Descriptions of new Oriental reptiles and batrachians. *Annals of the Magazine of Natural History*, ser. 6, 7:279–283.
- BOULENGER, G. A. 1894. On the herpetological fauna of Palawan and Balabac. *Annals of the Magazine of Natural History*, ser. 6, 6:81–90.
- BOULENGER, G. A. 1896. *Catalogue of the Snakes in the British Museum. III*. Taylor and Francis, London, UK. 727 pp.
- BOULENGER, G. A. 1897. Descriptions of new Malay frogs. *Annals and Magazine of Natural History*, ser. 5, 19:106–108.
- BROWN, R. M. 2007. *Introduction to Robert F. Inger's Systematics and Zoogeography of Philippine Amphibia*. Invited forward to the reprint of Inger's 1954 monograph. Pages 1–17 in *Systematics and Zoogeography of Philippine Amphibia*. Natural History Publications, Kota Kinabalu.
- BROWN, R. M. 2015. A new species of stream frog of the genus *Hylarana* from the mountains of southern Mindanao Island, Philippines. *Herpetologica* 71:223–233.
- BROWN, R. M., AND A. C. DIESMOS. 2009. Philippines, biology. Pages 723–732 in R. Gillespie and D. Clague, eds., *Encyclopedia of Islands*. University of California Press, Berkeley, California, USA.
- BROWN, R. M., A. C. DIESMOS, AND A. C. ALCALA. 2001 (2002). Application of lineage-based species concepts to oceanic frog populations: The effects of differing taxonomic philosophies on the estimation of Philippine biodiversity. *Silliman Journal* 42:133–162.
- BROWN, R. M., A. C. DIESMOS, M. B. SANGUILA, C. D. SILER, M. L. D. DIESMOS, AND A. C. ALCALA. 2012. Amphibian conservation in the Philippines. *FrogLog* 104:40–43.
- BROWN, R. M., AND J. C. GONZALEZ. 2007. A new forest frog of the genus *Platymantis* (Amphibia: Anura: Ranidae) from the Bicol Peninsula of Luzon Island, Philippines. *Copeia* 2007:251–266.
- BROWN, R. M., AND S. I. GUTTMAN. 2002. Phylogenetic systematics of the *Rana signata* complex of Philippine and Bornean stream frogs: reconsideration of Huxley's modification of Wallace's Line at the Oriental–Australian faunal zone interface. *Biological Journal of the Linnean Society London* 76: 393–461.
- BROWN, R. M., L. A. DE LAYOLA, A. LORENZO II, M. L. L. DIESMOS, AND A. C. DIESMOS. 2015. A new species of limestone karst inhabiting forest frog, genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae: subgenus *Lupacolus*) from southern Luzon Island, Philippines. *Zootaxa* 4048:191–210.
- BROWN, R. M., J. A. MCGUIRE, AND A. C. DIESMOS. 2000. Status of some Philippine frogs referred to *Rana everetti* (Anura: Ranidae), description of a new species, and resurrection of *Rana igorata* Taylor 1922.

Herpetologica 56:81–104.

- BROWN, R. M., AND C. D. SILER. 2013. Spotted stream frog diversification at the Australasian faunal zone interface, mainland versus island comparisons, and a test of the Philippine ‘dual-umbilicus’ hypothesis. *Journal of Biogeography* 41:182–195.
- BROWN, R. M., C. D. SILER, A. C. DIESMOS, AND A. C. ALCALA. 2009. The Philippines frogs of the genus *Leptobrachium* (Anura; Megophryidae) : taxonomic revision, phylogeny-based species delimitation, and descriptions of three new species. *Herpetological Monographs* 23:1–44.
- BROWN, R. M., C. D. SILER, C. H. OLIVEROS, J. A. ESSELSTYN, A. C. DIESMOS, P. A. HOSNER, C. W. LINKEM, A. J. BARLEY, J. R. OAKS, M. B. SANGUILA, L. J. WELTON, D. S. BLACKBURN, R. G. MOYLE, A. T. PETERSON, AND A. C. ALCALA. 2013. Evolutionary processes of diversification in a model island archipelago. *Annual Review of Ecology, Evolution, and Systematics* 44:411–435.
- BROWN, R. M., C. D. SILER, S. RICHARDS, A. C. DIESMOS, AND D. C. CANNATELLA. 2015. Multilocus phylogeny and a new classification for Southeast Asian and Melanesian forest frogs (family Ceratobatrachidae). *Zoological Journal of the Linnean Society* 174:130–168.
- BROWN, R. M., AND B. L. STUART. 2012. Patterns of biodiversity discovery through time: an historical analysis of amphibian species discoveries in the Southeast Asian mainland and adjacent island archipelagos. Pages 348–389 in Gower, D. J., K. G. Johnson, J. E. Richardson, B. R. Rosen, L. Rüber, and S. T. Williams, eds., *Biotic Evolution and Environmental Change in Southeast Asia*. Cambridge University Press, Cambridge, UK.
- BROWN, W. C., AND A. C. ALCALA. 1967. A new frog of the genus *Oreophryne* and a list of amphibians from Camiguin Island, Philippines. *Proceedings of the Biological Society of Washington* 80:65–68.
- BROWN, W. C., AND A. C. ALCALA. 1970a. A new species of the genus *Platymantis* (Ranidae) with a list of amphibians known from South Gigante Island, Philippines. *Occasional Papers of the California Academy of Sciences* 84:1–8.
- BROWN, W. C., AND A. C. ALCALA. 1970b. The zoogeography of the herpetofauna of the Philippine Islands, a fringing archipelago. *Proceedings of the California Academy of Sciences*, ser. 4, 38:105–130.
- BROWN, W. C., AND A. C. ALCALA. 1974. New frogs of the genus *Platymantis* (Ranidae) from the Philippines. *Occasional Papers of the California Academy of Sciences* 113:1–12.
- BROWN, W. C., AND A. C. ALCALA. 1977. A new frog of the genus *Rana* from the Philippines. *Proceedings of the Biological Society of Washington* 90:669–675.
- BROWN, W. C., AND A. C. ALCALA. 1982. A new cave *Platymantis* (Amphibia: Ranidae) from the Philippine Islands. *Proceedings of the Biological Society of Washington* 95:386–391.
- BROWN, W. C., AND A. C. ALCALA. 1994. Philippine frogs of the family Rhacophoridae. *Proceedings of the California Academy of Sciences*, ser. 4, 48:185–220.
- BROWN, W. C., A. C. ALCALA, A. C. DIESMOS, AND E. ALCALA. 1997a. Species of the *guentheri* group of *Platymantis* (Amphibia: Ranidae) from the Philippines, with descriptions of four new species. *Proceedings of the California Academy of Sciences*, ser. 4, 50:1–20.
- BROWN, W. C., R. M. BROWN, AND A. C. ALCALA. 1997b. Species of the *hazela* group of *Platymantis* (Amphibia: Ranidae) from the Philippines, with descriptions of two new species. *Proceedings of the California Academy of Sciences*, ser. 4, 49:405–421.
- BROWN, W. C., R. M. BROWN, A. C. ALCALA, AND R. FROST. 1997c. Replacement name for *Platymantis reticulatus* Brown, Brown, and Alcala, 1997 (Ranidae: Raninae). *Herpetological Review* 28:131.
- BROWN, W. C., A. C. ALCALA, AND A. C. DIESMOS. 1999a. Four new species of the genus *Platymantis* (Amphibia: Ranidae) from Luzon Island, Philippines. *Proceedings of the California Academy of Sciences*, ser. 4, 51:449–460.
- BROWN, W. C., A. C. ALCALA, P. S. ONG, AND A. C. DIESMOS. 1999b. A new species of *Platymantis* (Amphibia: Ranidae) from the Sierra Madre Mountains, Luzon Island, Philippines. *Proceedings of the Biological Society of Washington* 112:510–514.
- CIAT-CSI SRTM. 2015. *The CGIAR Consortium for Spatial Information*. <<http://srtm.csi.cgiar.org>>. Cited 1 October 2015.
- DIESMOS, A. C., R. M. BROWN, AND A. C. ALCALA. 2002. New species of narrow-mouthed frog (Amphibia: Anura: Microhylidae: genus *Kaloula*) from the mountains of southern Luzon and Polillo Islands,

- Philippines. *Copeia* 2002:1037–1051.
- DIESMOS, M. L. D., A. C. DIESMOS, C. D. SILER, V. T. VREDENBURG, AND R. M. BROWN. 2012. Detecting the distribution of the chytrid fungus in the Philippines. *FrogLog* 104:48–49.
- DIESMOS, A. C., R. M. BROWN, A. C. ALCALA, R. V. SISON, L. E. AFUANG, AND G. V. A. GEE. 2002. Philippine amphibians and reptiles. Pages 26–44 in P. S. Ong, L. E. Afuang, and R. G. Rosell-Ambal, eds., *Philippine Biodiversity Conservation Priorities: a Second Iteration of the National Biodiversity Strategy and Action Plan*. Department of the Environment and Natural Resources–Protected Areas and Wildlife Bureau, Conservation International Philippines, Biodiversity Conservation Program–University of the Philippines Center for Integrative and Developmental Studies, and Foundation for the Philippine Environment, Quezon City, Philippines.
- DIESMOS, A. C., M. L. DIESMOS, AND R. M. BROWN. 2006. Status and distribution of alien invasive frogs in the Philippines. *Journal of Environmental Science and Management, Philippines* 9:41–53.
- DIESMOS, A. C., A. C. ALCALA, C. D. SILER, AND R. M. BROWN. 2014. Status and conservation of Philippine amphibians. Pages 310–336 in H. Heatwole, H. and I. Das, eds., *Conservation Biology of Amphibians of Asia. Status and Decline of Amphibians: Eastern Hemisphere*. Natural History Publications (Borneo), Kota Kinabalu, Malaysia.
- DIESMOS, A. C., AND R. M. BROWN. 2011. Diversity, biogeography, and conservation of Philippine amphibians. Pages 26–49 in I. Das, A. Haas and A. A. Tuen, eds., *Biology and Conservation of Tropical Asian Amphibians*. Proceedings of the Conference “Biology of the Amphibians in the Sunda Region, Southeast Asia.” Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia.
- DUMÉRIL, A. H. A. 1853. Mémoire sur les batraciens anoures, de la famille des hylaeformes ou rainettes, comprennent la description d’un genre nouveau et de onze espèces nouvelles. *Annales des Sciences Naturelles, Zoologie et Biologie Animale, Paris, Serie 3*, 19:135–179.
- DUMÉRIL, A. M. C., AND G. BIBRON. 1841. *Erpétologie Générale on Histoire Naturelle Complète des Reptiles. Vol.8*. Librairie Encyclopedique de Roret, Paris, France. 24 pp.
- COPE, E. D. 1862. On some new and little known American Anura. *Proceedings of the Academy of Natural Sciences of Philadelphia* 14:151–159.
- FROST, D. R. 2015. *Amphibian Species of the World: an Online Reference. Version 6.0*. American Museum of Natural History, New York, USA. <<http://research.amnh.org/herpetology/amphibia/index.html>>. Cited 21 September 2015.
- FUITEN, A. M., L. J. WELTON, A. C. DIESMOS, A. J. BARLEY, B. OBERHEIDE, M. V. DUYA, E. L. B. RICO, AND R. M. BROWN. 2011. A new species of stream frog (*Sanguirana*) from the mountains of Luzon Island, Philippines. *Herpetologica* 67:89–103.
- GRAVENHORST, J. L. C. 1829. *Deliciae Musei Zoologici Vratislaviensis. Fasciculus primus. Chelonios et Batrachia*. Leopold Voss, Leipzig, Germany. 104 pp.
- GÜNTHER, A. 1873. Notes on some reptiles and batrachians obtained by Dr. Bernhard Meyer in Celebes and the Philippine Islands. *Proceedings of the Zoological Society of London* 1873:165–172.
- GÜNTHER, A. C. L. G. 1858. Neue Batrachier in der Sammlung des Britischen Museums. *Archiv für Naturgeschichte* 24:319–332.
- GÜNTHER, A. C. L. G. 1895. The reptiles and batrachians of the Natuna Islands. *Novitates Zoologicae* 2:499–502.
- INGER, R. F. 1954. Systematics and zoogeography of Philippine Amphibia. *Fieldiana Zoology* 33:183–531.
- INGER, R. F., AND K. J. FROGNER. 1979. New species of narrow-mouth frogs (genus *Microhyla*) from Borneo. *Sarawak Museum Journal* 27:311–322.
- KUHL, H., AND J. C. VAN HASSELT. 1822. Uittreksels uit breieven van de Heeren Kuhl en van Hasselt, aan de Heeren C. J. Temminck, Th. van Swinderen en W. de Haan. *Algemeene Konst–en Letter–Bode* 7:99–104.
- LINNAEUS, C. 1758. *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. 10th Edition. Volume 1*. Laurentii Salvii, Stockholm, Sweden. 824 pp.
- MOCQUARD, F. 1890. Recherches sur la faune herpetologique des îles de Bornéo et de Palawan. *Nouvelles Archives du Muséum d’Histoire Naturelle. Série 3. Paris* 2:115–168.

- MOCQUARD, F. 1892. Description de deux ophidiens et d'un batracien d'espèces nouvelles. *Le Naturaliste, Série 2, Paris* 6:35.
- OLSON, C. A., A. C. DIESMOS, AND K. H. BEARD. 2014. *Eleutherodactylus planirostris* (Greenhouse Frog) on Mindanao, Philippines. *Herpetological Review* 45:652–653.
- PETERS, W. C. H. 1863. Fernere Mittheilungen über neue Batrachier. *Monatsberichte der Königlichen Preussische Akademie des Wissenschaften zu Berlin* 1863:445–470.
- PETERS, W. C. H. 1867. Herpetologische Notizen. *Monatsberichte der Königlichen Preussische Akademie des Wissenschaften zu Berlin* 1867:13–37.
- PETERS, W. C. H. 1871. Über neue Reptilien aus Ostafrika und Sarawak (Borneo), vorzüglich aus der Sammlung des Hrn. Marquis J. Doria zu Genua. *Monatsberichte der Königlichen Preussische Akademie des Wissenschaften zu Berlin* 1871:566–581.
- REUTER H. I., A. NELSON, AND A. JARVIS. 2007. An evaluation of void filling interpolation methods for SRTM data. *International Journal of Geographic Information Science* 21:983–1008.
- ROSS, C. A., AND P. C. GONZALES. 1992. Amphibians and reptiles of Catanduanes Island, Philippines. *National Museum Papers, Manila* 2:50–76.
- ROWLEY, J., R. BROWN, R. BAIN, M. KUSRINI, R. INGER, B. STUART, G. WOGAN, N. THY, T. CHANARD, C. T. TRUNG, A. DIESMOS, D. T. ISKANDAR, M. LAU, L.T. MING, S. MAKCHAI, N. Q. TRUONG AND S. PHIMMACHAK. 2010. Impending conservation crisis for Southeast Asian amphibians. *Biology Letters* 6:336–338.
- SABAJ PEREZ, M. H. 2014. Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an Online Reference. Version 5.0 (28 June 2014). American Society of Ichthyologists and Herpetologists, Washington, DC. Available from: <http://www.asih.org/> (Accessed 1 June 2015).
- SCHLEGEL, H. 1837. *Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworfen, herausgegeben und mit einem erläuternden Texte begleitet. Part 1.* Arnz & Co., Düsseldorf, Germany. 141 pp.
- SCHNEIDER, J. G. 1799. *Historia Amphibiorum Naturalis et Literariae. Fasciculus Primus. Continens Ranas, Calamitas, Bufones, Salamandras et Hydros in Genera et Species Descriptos Notisque suis Distinctos.* Friederici Frommanni, Jena, Austria.
- SHAW, G. 1802. *General Zoology, or Systematic Natural History.* Vol.3. G. Kearsley, Thomas Davison, London, England, UK. 303 pp.
- SILER, C. D., C. W. LINKEM, A. C. DIESMOS, AND A. C. ALCALA. 2007. A new species of *Platymantis* (Amphibia: Anura: Ranidae) from Panay Island, Philippines. *Herpetologica* 63:351–364.
- SILER, C. D., A. C. ALCALA, A. C. DIESMOS, AND R. M. BROWN. 2009a. A new species of limestone forest frog, genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae) from eastern Samar Island, Philippines. *Herpetologica* 65:92–104.
- SILER, C. D., J. D. MCVAY, A. C. DIESMOS, AND R. M. BROWN. 2009b. A new species of fanged frog, genus *Limnonectes* (Amphibia: Anura: Dicroglossidae) from Southeast Mindanao Island, Philippines. *Herpetologica* 65:105–114.
- SILER, C. D., A. C. DIESMOS, C. W. LINKEM, M. L. DIESMOS, AND R. M. BROWN. 2010. A new species of limestone-forest frog, genus *Platymantis* (Amphibia: Anura: Ceratobatrachidae) from central Luzon Island, Philippines. *Zootaxa* 2482:49–63.
- SODHI, N. S., L. P. KOH, B.W. BROOK, AND P. K. L NG. 2004. Southeast Asian biodiversity: an impending disaster. *Trends in Ecology and Evolution* 19:654–660.
- STEJNEGER, L. 1905. Three new frogs and one new gecko from the Philippine Islands. *Proceedings of the United States National Museum* 28:343–348.
- STEJNEGER, L. 1908. Two new species of toads from the Philippines. *Proceedings of the United States National Museum* 33:573–576.
- STEJNEGER, L. 1910. Description of a new frog from the Philippine Islands. *Smithsonian Miscellaneous Collections* 52:437–439.
- STOLICZKA, F. 1870. Observations on some Indian and Malayan amphibia and reptilia. *Proceedings of the Asiatic Society of Bengal* 1870:105–109.
- SY, E. Y., J. C. MARTYR, AND A. C. DIESMOS. 2015. *Eleutherodactylus planirostris* (Greenhouse Frog) on

- Luzon, Philippines. *Herpetological Review* 46:56.
- SY, E. Y., D. VILLA-REAL, AND G. C. GAMOLO. 2014. *Kaloula pulchra* (Asiatic Painted Frog) on Cebu, Philippines. *Herpetological Review* 45:276–277.
- TAYLOR, E. H. 1920. Philippine Amphibia. *Philippine Journal of Science* 16:213–359.
- TAYLOR, E. H. 1922. Additions to the herpetological fauna of the Philippine Islands, I. *Philippine Journal of Science* 21:161–206.
- TAYLOR, E. H. 1923. Additions to the herpetological fauna of the Philippine Islands, III. *Philippine Journal of Science* 22:515–557.
- TAYLOR, E. H. 1960. On the caecilian species *Ichthyophis glutinosus* and *Ichthyophis monochrous*, with description of related species. *University of Kansas Science Bulletin* 40:37–120.
- TAYLOR, E. H., AND G. K. NOBLE. 1924. A new genus of discoglossid frogs from the Philippine Islands. *American Museum Novitates* 121:1–4.
- VAN OORT, P., AND S. MÜLLER. 1836. Aantekengingen gehouden op eene Reize over een Gedeelte van het Eiland Java, door de Leden der Natuurkundige Commissie. *Verhandelingen Bataviaasch Genootschap van Kunsten en Wetenschappen* 16:81–156.
- VON TSCHUDI, J. J. 1838. *Classification der Batrachier, mit berucksichtigung der fossilen Thiere dieser Abtheilung der Reptilien*. Petitpierre, Neuchâtel. 99 pp.
- WIEGMANN, A. F. A. 1834. Amphibien. Pages 433–522 in F. J. F. Meyen, ed., *Reise um die Erde ausgeführt auf dem Königlich Preussischen Seehandlungs-Schiffe Prinzes Louise, comandiert von Captain W. Wendt, in den Jahren 1830, 1831 und 1832 von Dr. F. J. F. Meyen. Dritter Theil. Zoologischer Bericht*. Sander'schen Buchhandlung (C. W. Eichhoff), Berlin, Germany.

AUTHOR CONTRIBUTIONS

CDS, RMB, ACD, and JLW conceived the ideas; ACD, NAH, ACA, RIC, LEA, GGD, RVS, MBS, MLD, EYS, LJW, RMB, and CDS participated in fieldwork; JLW, MBS, MLP, MJL, CSD, EAL, and CDS compiled and analyzed the dataset; NAH created the maps; DRD created the photo plates; JLW, NAH, DRD, RMB, and CDS revised analyzed the data; ACD and JLW led the writing; JLW, NAH, and CDS wrote the introduction and discussion; and JLW, CDS, NAH, DRD, and RMB edited drafts of the manuscript.

Distribution Maps and Photographs
Figures 2–44



FIGURE 2. Topographic map of the Philippine archipelago, with island names provided for larger islands. Numeric labels for smaller islands correspond to inset key.




FIGURE 3. Geographic range maps for members of the families Bombinatoridae (*Barbourula busuangensis*), and Bufonidae (*Ansonia mcgregori*, *A. muelleri*, and *Ingerophrynus philippinus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 4. Geographic range maps for members of the family Bufonidae (*Pelophryne albotaeniata*, *P. brevipes*, *P. lighti*, and *Rhinella marina*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 5. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis banahao*, *P. bayani*, *P. biak*, and *P. cagayanensis*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 6. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis cornutus*, *P. corrugatus*, *P. diesmosi*, and *P. dorsalis*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 7. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis guentheri*, *P. hazelae*, *P. indepressus*, and *P. insulatus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 8. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis isarog*, *P. lawtoni*, *P. levigatus*, and *P. luzonensis*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 9. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis mimulus*, *P. montanus*, *P. naomii*, and *P. negrosensis*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 10. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis paengi*, *P. panayensis*, *P. polillensis*, and *P. pseudodorsalis*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 11. Geographic range maps for members of the family Ceratobatrachidae (*Platymantis pygmaeus*, *P. rabori*, *P. sierramadrensis*, and *P. spelaeus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 12. Geographic range maps for members of the families Ceratobatrachidae (*Platymantis subterrestris*, *P. taylori*, and *Alcalus mariae*) and Dicroglossidae (*Fejervarya moodiei*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 13. Geographic range maps for members of the family Dicroglossidae (*Fejervarya vittigera*, *Hoplobatrachus rugulosus*, *Limnonectes acanthi*, and *L. diuatus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 14. Geographic range maps for members of the family Dicroglossidae (*Limnonectes feneri*, *L. leytensis*, *L. macrocephalus*, and *L. magnus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 15. Geographic range maps for members of the family Dicroglossidae (*Limnonectes micrixalus*, *L. palavanensis*, *L. parvus*, and *L. visayanus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 16. Geographic range maps for members of the families Dicroglossidae (*Limnonectes woodworthi*, *Occidozyga diminutiva*, and *O. laevis*), and Eleutherodactylidae (*Eleutherodactylus planirostris*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 17. Geographic range maps for members of the family Megophryidae (*Leptobrachium lumadorum*, *L. mangyanorum*, *L. tagbanorum*, and *Megophrys ligayae*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 18. Geographic range maps for members of the families Megophryidae (*Megophrys stejnegeri*), and Microhylidae (*Chaperina fusca*, *Kalophrynus sinensis*, and *Kaloula baleata*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 19. Geographic range maps for members of the family Microhylidae (*Kaloula conjuncta*, *K. kalingensis*, *K. kokacii*, and *K. picta*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 20. Geographic range maps for members of the family Microhylidae (*Kaloula pulchra*, *K. rigida*, *K. walteri*, and *Microhyla petrigena*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 21. Geographic range maps for members of the families Microhylidae (*Oreophryne anulata* and *O. nana*), and Ranidae (*Amnirana nicobariensis* and *Hylarana erythraea*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 22. Geographic range maps for members of the family Ranidae (*Lithobates catesbeianus*, *Pulchrana grandocula*, *P. mangyanum*, and *P. melanomenta*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 23. Geographic range maps for members of the family Ranidae (*Pulchrana moellendorffi*, *P. similis*, *Sanguirana albotuberculata*, and *S. aurantipunctata*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 24. Geographic range maps for members of the family Ranidae (*Sanguirana everetti*, *S. igorota*, *S. luzonensis*, and *S. sanguinea*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 25. Geographic range maps for members of the families Ranidae (*Sanguirana tipanan*, *Staurois natator*, and *S. nubilus*), and Rhacophoridae (*Kurixalus appendiculatus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 26. Geographic range maps for members of the family Rhacophoridae (*Nyctixalus pictus*, *N. spinosus*, *Philautus acutirostris*, and *P. everetti*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 27. Geographic range maps for members of the family Rhacophoridae (*Philautus leitensis*, *P. longicrus*, *P. poecilus*, and *P. schmackeri*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 28. Geographic range maps for members of the family Rhacophoridae (*Philautus surdus*, *P. surrufus*, *P. worcesteri*, and *Polypedates leucomystax*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 29. Geographic range maps for members of the families Rhacophoridae (*Polypedates macrotis*, *Rhacophorus bimaculatus*, and *R. pardalis*), and Ichthyophiidae (*Ichthyophis glandulosus*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.




FIGURE 30. Geographic range maps for members of the family Ichthyophiidae (*Ichthyophis mindanaoensis* and *I. weberi*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.



FIGURE 31. Photographs in life of A) *Barbourula busuangensis* (Bombinatoridae), B) *Ansonia mcgregori* (Bufonidae), C) *Ansonia muelleri* (Bufonidae), D) *Ingerophrynus philippinus* (Bufonidae), E) *Pelophryne brevipes* (Bufonidae), F) *Pelophryne lighti* (Bufonidae), G) *Rhinella marina* (Bufonidae), and H) *Rhinella marina* (Bufonidae). Photographs copyright Rafe M. Brown (A), Janalee P. Caldwell (H), and Cameron D. Siler (B, C, D, E, F, G).



FIGURE 32. Photographs in life of A) *Platymantis banahao* (Ceratobatrachidae), B) *Platymantis bayani* (Ceratobatrachidae), C) *Platymantis biak* (Ceratobatrachidae), D) *Platymantis cagayanensis* (Ceratobatrachidae), E) *Platymantis cornutus* (Ceratobatrachidae), F) *Platymantis corrugatus* (Ceratobatrachidae), G) *Platymantis diesmosi* (Ceratobatrachidae), H) *Platymantis dorsalis* (Ceratobatrachidae). Photographs copyright Rafe M. Brown (A, D, E, G), Arvin C. Diesmos (C), and Cameron D. Siler (B, F, H).



FIGURE 33. Photographs in life of A) *Platymantis guentheri* (Ceratobatrachidae), B) *Platymantis hazelae* (Ceratobatrachidae), C) *Platymantis insulatus* (Ceratobatrachidae), D) *Platymantis isarog* (Ceratobatrachidae), E) *Platymantis lawtoni* (Ceratobatrachidae), F) *Platymantis levigatus* (Ceratobatrachidae), G) *Platymantis luzonensis* (Ceratobatrachidae), and H) *Platymantis montanus* (Ceratobatrachidae). Photographs copyright Rafe M. Brown (C, D, G, H), Jason Fernandez (E), and Cameron D. Siler (A, B, F).



FIGURE 34. Photographs in life of A) *Platymantis negrosensis* (Ceratobatrachidae), B) *Platymantis paengi* (Ceratobatrachidae), C) *Platymantis polillensis* (Ceratobatrachidae), D) *Platymantis pygmaeus* (Ceratobatrachidae), E) *Platymantis rabori* (Ceratobatrachidae), F) *Platymantis sierramadrensis* (Ceratobatrachidae), G) *Platymantis spelaeus* (Ceratobatrachidae), and H) *Platymantis subterrestris* (Ceratobatrachidae). Photographs copyright Rafe M. Brown (D, F, H) Cameron D. Siler (A, B, E, G), and Luke Welton (C).



FIGURE 35. Photographs in life of A) *Platymantis taylori* (Ceratobatrachidae), B) *Alcalus mariae* (Ceratobatrachidae), C) *Fejervarya moodiei* (Dicroglossidae), D) *Fejervarya vittigera* (Dicroglossidae), E) *Hoplobatrachus rugulosus* (Dicroglossidae), F) *Limnonectes acanthi* (Dicroglossidae), G) *Limnonectes leytenis* (Dicroglossidae), and H) *Limnonectes macrocephalus* (Dicroglossidae). Photographs copyright Rafe M. Brown (B, H), Arvin C. Diesmos (A, E), Cameron D. Siler (C, D, G), and Scott Travers (F).



FIGURE 36. Photographs in life of A) *Limnonectes magnus* (Dicroglossidae), B) *Limnonectes palavanensis* (Dicroglossidae), C) *Limnonectes parvus* (Dicroglossidae), D) *Limnonectes visayanus* (Dicroglossidae), E) *Limnonectes woodworthi* (Dicroglossidae), F) *Occidozyga diminutiva* (Dicroglossidae), G) *Occidozyga laevis* (Dicroglossidae), and H) *Leptobrachium lumadum* (Megophryidae). Photographs copyright Rafe M. Brown (A, B, C, D, F) and Cameron D. Siler (E, G).




FIGURE 37. Photographs in life of A) *Leptobrachium mangyanorum* (Megophryidae), B) *Leptobrachium tagbanorum* (Megophryidae), C) *Megophrys ligayae* (Megophryidae), D) *Megophrys stejnegeri* (Megophryidae), E) *Chaperina fusca* (Microhylidae), F) *Kalophrynus sinensis* (Microhylidae), G) *Kaloula conjuncta* (Microhylidae), and H) *Kaloula kalingensis* (Microhylidae). Photographs copyright Rafe M. Brown (A, B, C, E, F), Arvin C. Diesmos (D), Cameron D. Siler (H), and Scott Travers (G).



FIGURE 38. Photographs in life of A) *Kaloula picta* (Microhylidae), B) *Kaloula pulchra* (Microhylidae), C) *Kaloula rigida* (Microhylidae), D) *Kaloula walteri* (Microhylidae), E) *Microhyla petrigena* (Microhylidae), F) *Oreophryne anulata* (Microhylidae), G) *Oreophryne nana* (Microhylidae), and H) *Hylarana erythraea* (Ranidae). Photographs copyright Rafe M. Brown (C, D, F), Arvin C. Diesmos (E), Cameron D. Siler (A, G, H), and Scott Travers (B).



FIGURE 39. Photographs in life of A) *Lithobates catesbeianus* (Ranidae), B) *Pulchrana granocula* (Ranidae), C) *Pulchrana mangyanum* (Ranidae), D) *Pulchrana moellendorffi* (Ranidae), E) *Pulchrana similis* (Ranidae), F) *Sanguirana albotuberculata* (Ranidae), G) *Sanguirana aurantipunctata* (Ranidae), and H) *Sanguirana everetti* (Ranidae). Photographs copyright Rafe M. Brown (D, G), Janalee P. Caldwell (A), and Cameron D. Siler (B, C, E, F, H).



FIGURE 40. Photographs in life of A) *Sanguirana igorota* (Ranidae), B) *Sanguirana luzonensis* (Ranidae), C) *Sanguirana sanguinea* (Ranidae), D) *Sanguirana tipanan* (Ranidae), E) *Staurois natator* (Ranidae), F) *Staurois nubilus* (Ranidae), G) *Kurixalus appendiculatus* (Rhacophoridae), and H) *Nyctixalus spinosus* (Rhacophoridae). Photographs copyright Rafe M. Brown (A, C, D, F) and Cameron D. Siler (B, E, G, H).



FIGURE 41. Photographs in life of A) *Nyctixalus spinosus* (Rhacophoridae), B) *Philautus acutirostris* (Rhacophoridae), C) *Philautus everetti* (Rhacophoridae), D) *Philautus leitensis* (Rhacophoridae), E) *Philautus longicrus* (Rhacophoridae), F) *Philautus poecilus* (Rhacophoridae), G) *Philautus surdus* (Rhacophoridae), and H) *Philautus worcesteri* (Rhacophoridae). Photographs copyright Rafe M. Brown (B, C, E, F, H) and Cameron D. Siler (A, D, G).



FIGURE 42. Photographs in life of A) *Polypedates leucomystax* (Rhacophoridae), B) *Polypedates macrotis* (Rhacophoridae), C) *Rhacophorus bimaculatus* (Rhacophoridae), D) *Rhacophorus pardalis* (Rhacophoridae), E) *Ichthyophis glandulosus* (Ichthyophiidae), and F) *Ichthyophis glandulosus* (Ichthyophiidae). Photographs copyright Rafe M. Brown (B, E, F) and Cameron D. Siler (A, C, D).




FIGURE 43. Geographic range maps for members of the families Ceratobatrachidae (*Platymantis quezoni*) and Ranidae (*Pulchrana guttmani*). Points represent museum vouchered specimens with georeferenced locality information overlaid on a topographic map of the Philippines.



FIGURE 44. Photograph in life of (A) *Platymantis quezoni* (Ceratobatrachidae). Photograph copyright Rafe M. Brown.

Page intentionally left blank

Species of *Macromitrium* (Orthotrichaceae) New to the Mindanao Region and the Philippines with One Species New to Science

Benito C Tan ¹ and James R. Shevock ²

¹ University Herbarium, 1001 Valley Life Sciences, University of California, Berkeley, California 94720-2465 U.S.A. Email: btakakia@yahoo.com; ² California Academy of Sciences, Department of Botany, 55 Music Concourse Drive, Golden Gate Park, San Francisco, California 94118-4503 U.S.A. Email: jshevock@calacademy.org

Numerous collections of *Macromitrium* were made during a joint expedition between the California Academy of Sciences and the Central Mindanao State University in 2014 and 2015 from the region of Mindanao in the Philippines. Two *Macromitrium* species collected from the Mindanao Region are new records for the Philippine flora, along with a species, *M. eddyi*. Range extensions of four species of *Macromitrium* across the island of Mindanao are mentioned. A review of the species diversity of *Macromitrium* between the Luzon Island region and the Mindanao region is also provided.

KEYWORDS: Asian mosses, bryophyte inventory, Malesia, Mindanao Island, new records, Papua New Guinea, Philippines, species distribution

Mindanao Island (ca 104,630 km²) and its off shore, small islands, although slightly smaller than and ranking second in total land area next to Luzon Island (ca 109,965 km²), have only half the number of moss species documented for the latter (Linis and Tan 2008). However, recent reports of collecting activities of mosses of Mindanao Island have added 60 new species records to the island (Tan and Shevock, 2014; Azuelo et al. 2015; Tan et al. 2015). Here we focus our study on the epiphytic moss genus, *Macromitrium* Brid., which has a different assemblage of species occurring on these two major Philippine island groups.

Early in this century, 21 species of *Macromitrium* were documented from the Philippines with 18 species reported from Luzon and 13 species from Mindanao (see Tan and Iwatsuki 1991; Tan et al. 2000). The two new expeditions organized by the California Academy of Sciences and the CEBREM Office of Central Mindanao State University in 2014 and 2015 have added three new records of *Macromitrium* (Tan et al. 2015; this publication), and one reported below as a species new to science.

An updated list of the 19 species and one variety of *Macromitrium* in Mindanao is appended below based on Tan et al. (2000), Linis (2010), Tan et al. (2015) and this publication. Of these, *M. archboldii* E.B. Bartram, *M. eddyi* B.C. Tan & Shevock sp. nov., *M. mindorensis* Broth., *M. ochraceum* (Dozy & Molk.) Müll. Hal. and *M. tylostomum* Mitt. ex Bosch & Sande Lac. are not documented from Luzon Island. On the other hand, *M. benguetense* Williams, *M. fasciculare* Mitt., *M. formosae* Card., *M. nepalense* (Hook. & Grev.) Schwägr., and *M. robinsonii* Williams still have no specimens found in Mindanao. It is noteworthy that Luzon has received considerably more bryophyte inventory work since it is within the Manila area compared to distant Mindanao. We speculate that additional Wallacea-Australian *Macromitrium* are unlikely to occur in the northern

half of the Philippine archipelago based on this collection history. However, we expect additional *Macromitrium* species documented from either adjacent Indonesia or Papua New Guinea could be discovered in Mindanao with additional field inventory.

NEW SPECIES

Dr. Alan Eddy (1937–1998), while affiliated with the herbarium of the British Museum (BM), was working on a multi-volumed *Handbook of Malesian Mosses* up to the time of his death. The third and final fascicle he published includes the moss family Orthotrichaceae (Eddy 1996). *Macromitrium* is a very large and complex genus with over 350 species (Crosby et al. 2000). During the development of the *Macromitrium* treatment he encountered a single specimen that could not be assigned to any of the known Malesian species. He decided to include this specimen and reference it simply as '*Macromitrium* sp.' with the hope that other bryologists would eventually encounter it. Therefore, he was the first person to recognize this *Macromitrium* as likely to be new to science. Although he decided not to name it as a new species based on a single sample, he nonetheless provided a detailed description (p. 38) and prepared an illustration (fig. 350) of this taxon based on the specimen labeled as "*Stevens no. 55716*" (see Eddy 1996). While examining our recently collected *Macromitrium* collections from Mindanao, we used the *Macromitrium* key in Eddy (1996) and realized that one of our unknown specimens matched the illustration and description entry as '*Macromitrium* sp.' We borrowed this collection from BM and compared it to the Philippine specimen (*Shevock 44672*) from Mt. Kitanglad Range Natural Park. We determined the species to be one and the same.

Macromitrium eddyi B. C. Tan & Shevock, sp. nov.

Figures 1–3.

HOLOTYPE: **Papua New Guinea:** Milne Bay District, Raba-Raba Sub-district, bottom of scarp of Tantam Plateau, Mt. Suckling, 1645 m., in shaded forest, common on wood, 20 Jul 1972, coll. P.F. Stevens [*LAE 55716*] (BM!; isotypes, CANB, E, L, LAE). **PARATYPE:** **Philippines: Mindanao Island:** Bukidnon Province, on access dirt road to trail less than 0.5 km above Lantapan Village toward Mt. Dulang-Dulang in Kitanglad Range Natural Park, on trunk of *Gmelina* in disturbed forest near cultivated field, 20 Apr 2014, *Shevock 44672* (CAS; isoparatype, CMUH, NY, UC).

The new species can be identified easily using the key to the species of *Macromitrium* published in Eddy (1996). Morphologically the new species is identified by its slender and long branches with leaves arranged in five straight, longitudinal rows, when wet. Other diagnostic characters include leaves with acute to acuminate apices, percurrent to shortly excurrent costae, papillose upper leaf cells, and tuberculate lower leaf cells. In addition, the basal leaf cells are thick-walled with straight to curved lumina.

Below we reproduce verbatim the species description prepared by Eddy (1996) in whose honor we name this new species in recognition of his great contribution to our knowledge of Malesian mosses. Additionally, we include herewith photographs of the specimen of the Philippine paratype to complement the nice illustration of what is now *M. eddyi* that appears in Eddy (1996).

"Plants yellowish, slender, with elongate, sparingly ramifying branches up to 5 cm long. Leaves closely set, squarrose-recurved and markedly pentastichous when moist, erect and appressed with crisped upper limbs when dry, rendering the branches string-like in appearance; triangular lanceolate, canaliculate, finely acuminate, up to 3 mm long; apex finely acute, pellucid. Costa percurrent to short excurrent. Upper lamina cells small, isodiametric, ca 6–9 μ m diameter, densely pluripapillose and obscure; lower lamina cells elongate with strongly thickened walls and




FIGURE 1. *Macromitrium eddyi* B.C. Tan & Shevock from the holotype, *Stevens 55716* (BM). A. Leaf apex with percurrent costa of elongate cells and upper laminal cells multipapillose. B. Leaves. C. Lower laminal cells elongate with thickened walls and narrow, straight to somewhat curved lumina cells conspicuously tuberculate. Scale bars: A, C: 5 μ m, B: 0.5 mm.



FIGURE 2. Habit of *Macromitrium eddyi* stems in a dry state. *Shevock 44672* (CAS).



FIGURE 3. Growth habit of *Macromitrium eddyi* from *Shevock 44672* (CAS).

narrow, straight lumina, conspicuously tuberculate. Perichaetial leaves narrowly triangular-lanceolate, plicate, very finely acuminate and filiform-pointed. Seta short, about 3 mm long, smooth; capsule ovoid, smooth, small, the urn about 1 mm long and 0.8 mm wide; peristome of pale, rather fugacious, triangular exostome teeth; calyptra naked.”

NEW SPECIES RECORDS FOR THE PHILIPPINES

Macromitrium tylostomum Mitt. ex Bosch & Sande Lac.

SPECIMEN EXAMINED:— **Mindanao Island:** North Cotabato Province: Mt. Apo, trail to Plot H of EDC, ca 1900 m, 29 Apr 2014, *Tan 2014-132* (CAS, UC).

Among members of *Macromitrium* in the Philippines, this is one of three species that have the laminal cells rounded to quadrate in shape from apex to near base. It differs from the other two species, *M. orthostichum* Nees ex Schwägr. and *M. falciculatum* Müll. Hal. in having smooth and not papillose basal laminal cells. Although the upper and middle laminal cells are smooth, thin-walled and budging, the basal laminal cells of *M. tylostomum* are thick-walled and decorated with tall papillae. The cylindrical capsule attached to a short, smooth seta is also distinctive. Outside of the Philippines, *M. tylostomum* is known from Java, Sumatra and Papua New Guinea.

Macromitrium yuleanum Broth. & Geheeb

SPECIMENS EXAMINED:— **Mindanao Island:** Bukidnon Province: Mt. Kitanglad Range Natural Park, midslope of Mt. Dulan-Dulang Peak along the ridge trail separating headwaters of Alanib River and Magnao River, on fallen branch in mixed hardwood forest, 21–23 Apr 2014, *Shevock 44727, 44779, 44842* (CAS, CMUH, NY, UC).

According to Vitt et al. (1995), this is the most common, large sized and variable species of *Macromitrium* at high elevation in Huon Peninsula of Papua New Guinea and has been described as a new species at least seven times. The Mindanao collections cited above fit well with the species description presented in Vitt et al. (1995). They differ mainly in having widely erect-spreading (not squarrose-recurved) leaves when wet, more narrowly acuminate leaf apices, and a calyptra with noticeably stiff and long hair.

Macromitrium yuleanum is best identified by its narrowly ovate-lanceolate to oblong lanceolate leaves with percurrent to short, excurrent costa, and a weakly differentiated border of 1 to 2 rows of short rectangular, smooth cells near the denticulate apex and along the margins of the upper half of leaf. The laminal cells vary from mammillose to unipapillose. The setae are long, reaching more than 15 mm. The perichaetial leaves have a very long excurrent costa, nearly of the same length of lamina in some leaves.

Among the local congeners, *M. longicaule* Müll. Hal. is most similar in plant habit and leaf characters to *Macromitrium yuleanum*, but the former has percurrent leaf costa and short seta less than 5 mm long. Likewise, the somewhat look-alike *M. macrosporum* Broth. differs from *M. yuleanum* in having the percurrent leaf costa that ends at an incurved and entire apex. The leaf cells are nearly smooth and only slightly mammillose, but not unipapillose like in some leaves of *Macromitrium yuleanum*. The calyptra of *M. macrosporum* is also smooth with no hairs. *Macromitrium yuleanum* can also be mistaken for *M. ochraceum*, which is more commonly encountered in Mindanao Island. According to Eddy (1996), it can be separated from the latter by its less acute and shorter leaf apices with setae twice as long as that of *M. ochraceum*.

Macromitrium yuleanum was previously viewed as an endemic species of New Guinea and Solomon Islands.

RANGE EXTENSIONS AND NEW SPECIES RECORDS FOR THE MINDANAO REGION

***Macromitrium blumei* Nees ex Schwägr. [syn. *M. zollingeri* Dozy & Molk.; *M. blumei* var. *zollingeri* (Mitt. ex Bosch & Sande Lac.) S. L. Guo, B. C. Tan & Virtanen]**

SPECIMENS EXAMINED:— **Mindanao Island:** Bukidnon Province, saddle below the summit of Mt. Dulang-Dulang in Mt. Kitanglad Range Natural Park, on branches, 24 Apr 2014, *Shevock & B.C. Tan* 44874 (CAS, CMUH, UC); *ibid*, Mt. Kiamo, along ridge trail to summit, on fallen hardwood branch, 7 May 2014, *Shevock & B.C. Tan* 45141 (CAS, CMUH, UC); *ibid*, Mt. Limbawon, trail along ridge top at 1885 m elev. from campsite about 9 km distance above Kibalabag village, on branch of *Dacrydium* in open site of montane forest, 30 Jun 2015, *Shevock et al.* 47042 (CAS, CMUH, UC); Davao Oriental Province, along trail from Camp 2 near pygmy forest to summit of Mt. Hamiguitan, 1175–1350 m elev., on small diameter tree trunk in sun, 22 Jun 2015, *Shevock & Yorong* 46903, 46912 (CAS, CMUH, UC).

This is a common species in the Philippines and also a widespread member of the genus in Malesia. On Mindanao Island, it has been reported only from Mt. Candoon in Bukidnon Province and Mt. Apo Natural Park. The collections cited above expand the distribution of this species in Mindanao.

Macromitrium blumei is interpreted by us as a polymorphic species that exhibits both smooth and strongly bulging upper leaf cells. The synonymy between *M. blumei* and *M. zollingeri* was suggested by Eddy (1996) and agreed by us after we examined many packets of Mindanao specimens of these two taxa. The coiling leaves around the short branches, when dry, give the shoot a characteristic rope-like appearance (Eddy 1996). The small plant size, coupled with a somewhat asymmetrically obtuse leaf apex and a short to moderately long, excurrent costa, are additional diagnostic characters of this species. *Macromitrium blumei* is a widespread and common species in Malesia.

***Macromitrium cuspidatum* Hampe**

SPECIMENS EXAMINED:— **Camiguin Island:** Municipality of Mambajao, Mt. Hibok-Hibok, along Tagdo Trail to the summit at 710 m elev., on tree trunk in filtered sunlight, 8 Jul 2015, *Shevock & B.C. Tan* 47157 (CAS, CMUH, UC). **Mindanao Island:** Davao Oriental Province, along trail from Camp 2 near pygmy forest to summit of Mt. Hamiguitan, 1040 m elev., on tree trunk in filtered sunlight, 22 Jun 2015, *Shevock & Yorong* 46898 (CAS, CMUH, UC); *ibid*, along trail from Camp 2 to Twin Falls, mid-slope of Mt. Hamiguitan, 900–960 m elev., on tree trunk in filtered sunlight, 23 Jun 2015, *Shevock & Yorong* 46937, 46953 (CAS, CMUH, UC).

This is a distinctive species of *Macromitrium* in the Philippine moss flora and can be easily identified by the nearly all oblong to elongate and smooth leaf cells, coupled with a long excurrent costa. The plants are yellowish green in color and moderately large in size, which make the sporophyte look small in comparison. It has been reported from many places in island groups of Luzon, Mindoro, Palawan and Visayan Region (Tan and Iwatsuki 1991), but not from Mindanao Island until a report on the mosses of Mt. Kiamo in Bukidnon Province, was published by Tan et al. (2015). The above cited collections provide a north and south distribution range extension across Mindanao Island. *Macromitrium cuspidatum* is a widespread species in Malesia but not as commonly encountered as *M. blumei*.

***Macromitrium microstomum* (Hook. & Grev.) Schwägr. [syn. *M. reinwardtii* Schwägr.]**

SPECIMEN EXAMINED:— **Mindanao Island:** North Cotabato Province, Mt. Apo, toward Lake

Ma-ag above the Geothermal Production Field of EDC, 1 May 2014, *Shevock 45012* (CAS, CMUH, UC).

According to Eddy (1996) and Vitt et al. (1995), this species is characterized by having smooth and flat lamina cells throughout the leaf, ovoid capsules borne on a long seta reaching often to 1.5 cm, and a naked, plicate calyptra.

Macromitrium microstomum, had been known earlier from the Philippines by its synonym, *M. reinwardtii*, and had already been reported from Luzon, Mindoro, and some islands in the Visayan Region (Tan and Iwatsuki, 1991); we now add Mindanao to the list. Otherwise, *Macromitrium microstomum* is widespread throughout tropical SE Asia reaching Australia, New Zealand, and Oceania. It is known also from Mexico, Guatemala, Costa Rica and the Caribbean Islands (Vitt et al. 1995).

***Macromitrium ochraceum* (Dozy & Molk.) Müll. Hal.**

SPECIMENS EXAMINED:— **Mindanao Island:** Bukidnon Province, Mt. Kiamo, along ridge trail to summit in mixed hardwood-podocarp shrubland, on small tree trunk, 7 May 2014, *Shevock & B.C. Tan 45171* (CAS, CMUH, UC); ibid, Mt. Limbawon, trail along ridge top at 1800–1885 m elev. from campsite about 8–9 km distance above Kibalabag village, on ground and branch of *Dacrydium* in open site of montane forest along a cascading stream, 30 Jun 2015, *Shevock et al. 47037, 47041* (CAS, CMUH, UC); Davao Oriental Province, Mt. Hamiguitan, volcanic rock wall, 22 Jun 2015, *Shevock & Yorong 46919* (CAS, CMUH, UC), hardwood branch, 24 Jun 2015, *Shevock 46972* (CAS, CMUH).

Although this species is not new to Mindanao Island, it was previously known from only a single locality. These collections represent a distribution range extension of this large-sized *Macromitrium*. Bartram (1939) commented that “the long, ruddy, sparingly branched secondary stems and the very rough setae quickly establish the identity of *M. ochraceum*”. According to Vitt et al. (1995), its smooth to slightly bulging, upper laminal cells and the narrowly elongate, straight, thick-walled and tuberculate basal leaf cells add to its species distinctiveness. A good illustration of this species appears in Eddy (1996).

On Mindanao, *M. ochraceum* is rather variable in its plant size, and, thus, also the coarseness of its stems. In herbarium collections, specimens of *M. ochraceum* are likely to be misidentified as *M. longicaule* Müll. Hal., which has a similarly large plant size, long branches, and curvy, dry leaf foliation. The former species however, has short and papillose setae less than 1 cm long, whereas the latter has smooth setae more than 1 cm long. The leaf cells of *M. ochraceum* are smooth to mamillate and bulging, not pluripapillose like the leaf cells of *M. longicaule*. *Macromitrium ochraceum* is a widespread species at higher elevations in Malesia.

ACKNOWLEDGMENTS

We are indebted to the administration of the California Academy of Sciences for funding support given to the 2014 and 2015 CAS-CMUH botanical expeditions to Mindanao. The organizational leadership and field assistance of Prof. Victor Amoroso at CEBREM of the Central Mindanao University (CMUH) and his office staff, especially Mr. Fulgent Coritico, are much appreciated. Collecting permits were authorized by the offices of the Philippines Department of the Environment and Natural Resources (DENR). The authors acknowledge the use of photographic and microscopic facilities at the Howell Lab, Department of Botany, California Academy of Sciences, and we especially thank Wen-zhang Ma for design of the *Macromitrium eddyi* microscopic plate as Fig. 1. The habit images of *Macromitrium eddyi* were created by the GigaMacro Magnify 2

imaging system, Project Lab, California Academy of Sciences with credit given to the photographer Kathryn Whitney. Also, we again want to take note, with thankful appreciation, for the loan of Dr. Eddy's material labeled '*Macromitrium* sp.' by the British Museum (BM). Lastly, we gratefully acknowledge the thoughtful comments provided by Dr. Thomas Daniel who perused an early version of the manuscript and to two anonymous reviewers.

LITERATURE CITED

- AZUELO, A., G. B. C. TAN, J. R. SHEVOCK, A. MANUEL, A. YORONG, AND L. G. SARIANA. 2015. Mosses new for Mindanao Island, Republic of the Philippines. *Proceedings of the California Academy of Sciences*, ser. 4, 62:127–134.
- BARTRAM, E. B. 1939. Mosses of the Philippines. *Philippine Journal of Science* 68:1–423.
- CROSBY, M. R., R. E. MAGILL, AND S. HE. 2000. *A Checklist of the Mosses*. Missouri Botanical Garden, Saint Louis, Missouri, USA. [iv] + 320 pp.
- EDDY, A. 1996. *A Handbook of Malesian Mosses, Vol. 3 — Splachnobryaceae to Leptostomataceae*. The History Museum (Natural History), London, England, UK.
- LINIS, V. C. 2010. The moss flora of Camiguin Island, Philippines and its floristic relations to some adjacent islands in the archipelago. *Telopea* 12:525–542.
- LINIS, V. C., AND B. C. TAN. 2008. Progress of studies on phytogeography and biodiversity of Philippine moss flora from 1991 to 2006. Pages 13–22 in H. Mohamed, B. B. Baki, A. Nasrulhaq-Boyce, and P. K. Y. Lee, eds., *Bryology in the New Millennium*. University of Malaya, Kuala Lumpur.
- TAN, B. C., AND Z. IWATSUKI. 1991. A new annotated Philippine moss checklist. *Harvard Papers in Botany* 3:1–64.
- TAN, B. C., L. LUBOS, AND U. SCHWARZ. 2000. New and biogeographically noteworthy records of Philippine mosses from Mindanao Island. *Tropical Bryology* 18:27–37.
- TAN, B. C., AND J. R. SHEVOCK. 2014. Noteworthy mosses collected from the 2014 joint expedition of CAS and CMU new to Mindanao Island of the Philippines. *Bryophyte Diversity and Evolution* 36:22–30.
- TAN, B. C., J. R. SHEVOCK, V. AMOROSO, AND F. CORITICO. 2015. Mosses new for Mindanao Island, The Philippines III. *Bulletin Natural Science Museum Tokyo Ser. B*. 41:91–97.
- VITT, D. H., T. KOPONEN, AND D. H. NORRIS. 1995. Bryophyte flora of the Huon Peninsula, Papua New Guinea. LV. *Desmotheca*, *Groutiella*, *Macrocoma* and *Macromitrium* (Orthotrichaceae, Musci). *Acta Botanica Fennica* 154:1–94.

Appendix I

List of 19 *Macromitrium* species and one variety reported for the Mindanao Region. An asterisk (*) indicates a new distribution record for a species reported in this publication, while two asterisks (**) indicate a species new to science.

- Macromitrium angustifolium* Dozy & Molk.— Camiguin Island: Mt. Timpoong, Mt. Hibok-Hibok (Linis, 2010).
- Macromitrium archboldii* E.B. Bartram — Mindanao Island: Mt. Kitanglad Natural Park (Tan et al. 2000).
- Macromitrium blumei* Nees ex Schwägr.— Mindanao Island: Mt. Candoon, Mt. Apo, Mt. Kitanglad Natural Park, Mt. Kiamo, Mt. Limbawon, Mt. Hamiguitan (Bartram, 1939; Tan and Iwatsuki, 1991; this publication).
- Macromitrium cuspidatum* Hampe — Mindanao Island: Mt. Kiamo, Mt. Hamiguitan (Tan et al. 2015); (this publication).
- ***Macromitrium eddyi* B.C. Tan & Shevock, sp. nov.— Mindanao Island: Mt. Kitanglad Natural Park (this publication).
- Macromitrium falciculatum* Müll. Hal.— Mindanao Island: Banga in Zamboanga Province (Bartram, 1939; Tan and Iwatsuki, 1991).
- Macromitrium foxworthyi* Broth.— Mindanao Island: Sax River in Zamboanga Province (Tan and Iwatsuki, 1991).
- Macromitrium fuscescens* Schwägr. [syn. *M. semipellucidum* Dozy & Molk.] — Mindanao Island: Camp Keithley in Lake Lanao Province, Agusan Province (Bartram, 1939; Tan and Iwatsuki, 1991).
- Macromitrium incurvifolium* (Hook. & Grev.) Schwägr. [syn. *M. subtile* Schwägr., *M. subuligerum* Bosch & Sande Lac.] — Camiguin Island: Mt. Timpoong, Mt. Hibok-Hibok. Mindanao Island: Alag River (Tan and Iwatsuki, 1991; Linis, 2010).
- Macromitrium longicaule* Müll. Hal.— Mindanao Island: Mt. Apo, Mt. Kitanglad (Tan and Iwatsuki, 1991).
- Macromitrium macrosporum* Broth. [syn. *Macromitrium goniostomum* Broth.] — Mindanao Island: Mt. Apo (Bartram, 1939; Tan and Iwatsuki, 1991).
- **Macromitrium microstomum* (Hook. & Grev.) Schwägr. [syn. *M. reinwardtii* Schwägr.] — Mindanao Island: Mt. Apo (this publication).
- Macromitrium mindorense* Broth.— Mindanao Island: Davao Province (Tan and Iwatsuki, 1991).
- Macromitrium ochraceum* (Dozy & Molk.) Müll. Hal.— Mindanao Island: Mt. Malindang, Mt. Kiamo, Mt. Limbawon (Bartram, 1939; Tan and Iwatsuki, 1991; this publication).
- Macromitrium orthostichum* Nees ex Schwägr.— Camiguin Island: Mt. Timpoong, Mt. Hibok-Hibok. Mindanao Island: Sax River in Zamboanga Province (Bartram, 1939; Tan and Iwatsuki, 1991; Linis, 2010).
- Macromitrium salakanum* Müll. Hal.— Camiguin Island: Mt. Timpoong, Mt. Hibok-Hibok (Bartram, 1939; Tan and Iwatsuki, 1991; Linis, 2010).
- Macromitrium salakanum* Müll. Hal. ssp. *celebense* (Paris) M. Fleisch.— Mindanao Island: Mt. Apo, Sax River in Zamboanga Province, Agusan Province (Bartram, 1939; Tan and Iwatsuki, 1991).
- Macromitrium sulcatum* (Hooker) Bridel — Mindanao Island: Mt. Candoon (Bartram, 1939; Tan and Iwatsuki, 1991).
- **Macromitrium tylostomum* Mitt. ex Bosch & Sande Lac.— Mindanao Island: Mt. Apo (this publication).
- **Macromitrium yuleanum* Broth. & Geh.— Mindanao Island: Mt. Kitanglad Range Natural Park (this publication).

Page intentionally left blank

Index to Taxonomic Names and Major Geographic Regions

A

Africa 451
 Madagascar 451, 453, 454, 455
Alcalus 461, 508, 531
 mariae 461, 508, 531
Amnirana 477, 517
 nicobariensis 477, 517
Anisotes 451, 452, 453, 454, 455
 divaricatus 454
 madagascariensis 454
 perplexus 454
 subcoriaceus 454
 tablensis **sp. nov.** 451, 452, 453, 454, 455
 venosus 454
Ansonia 459, 499, 527
 mcgregori 459, 499, 527
 muelleri 459, 499, 527
Anura 459, 491, 492, 493, 494
Asia (eastern) 469, 476;
 China (see under main heading)
Asia (Southeast) 457, 541
 Indonesia (see under main heading)
 Philippine Islands (see under main heading)
 Malaysia (see under main heading)
 Malesia (see under main heading)
Asia (Southwest) 451
 Arabian Peninsula 451
Auletris 477
 bilineatus 477
Australia 547

B

Barbourula busuangensis 459, 499, 527
Bombinator 460, 474
 baleatus 474
 horridus 460
 Hylaedactylus (as subgenus) *baleatus* 474
Bombinatoridae 459, 489, 499, 527
Bufo 459, 460
 agua 460
 albicans 460
 angustipes 460
 biporcatus philippinicus 459
 brasiliensis 460
 brevipes 460
 divergens 459
 horridus 460
 humeralis 460
 lazarus 460

maculiventris 460
 marinis 460
 marinus 460
 fluminensis 460
 horribilis 460
 napensis 460
 mcgregori 459
 muelleri 459
 philippinicus 459
 pithecodactylus 460
 pythecodactylus 460
Bufonidae 459, 489, 499, 500, 527

C

Calamita 477
 bilineatus 477
Callula 474, 475
 baleata 474
 Kallula (as subgenus) *pulchra* 475
 macrodactyla 475
 picta 475
 pulchra 475
Calohyla 474, 475
 celebensis 474
 pulchra 475
Calophrynus 474
 pleurostigma var. *sinensis* 474
Caloula 475
 pulchra 475
Canada 445
 British Columbia 445
 Vancouver Island 445
Carabidae 441
Caribbean Islands 472, 547
 Bahamas 472
 New Providence Island 472
Central America 547
 Costa Rica 547
 Guatemala 547
 Mexico (see under main heading)
Ceratobatrachidae 461, 489, 501, 502, 503, 504, 505,
 506, 507, 508, 528, 529, 530, 531, 539
Chalcorana 480, 481, 482
 albotuberculata 480
 everetti 481
 luzonensis 481
 tipanan 482
Chaperina 474, 476, 514, 533
 beyeri 474

fusca 474, 514, 533

visaya 476

Chaunus 461

marinus 461

China 469, 476

Coleoptera 441

Comoros Archipelago 451

Cornifer 465

guentheri 465

Cornufer 462, 463, 465, 466, 467, 486

cornutus 462

dorsalis 463

guentheri 463

hazela 463

ingeri 463

jagorii 463

laticeps 463

meyeri 463

montanus 465

polillensis 466

rivularis 463

subterrestris 467

worcesteri 463

D

Dicroglossidae 468, 489, 508, 509, 510, 511, 512, 531, 532

Dicroglossus 468

cancrivorus 468

Docidophryne 460

agua 460

Lazarus 460

E

Edwardtayloria 484

picta 484

spinosa 484

Eleutherodactylidae 472, 489, 512

Eleutherodactylus 472, 489, 512

Euhyas (as subgenus) *planirostris* 472

planirostris 472, 489, 512

planirostris 472

ricordii planirostris 472

Euhyas 472

planirostris 472

Euphlyctis 468, 469, 470, 471

cancrivora 468

limnocharis vittigera 468

magna 470

macrocephala 470

microdisca parva 471

palavanensis 471

tigerina rugulosa 469

F

Fejervarya 468, 469, 508, 509, 531

cancrivora 468

moodiei 468, 508, 531

raja 468

vittigera 469

vittigera 468, 509, 531

G

Gymnophiona 488

H

Halophila 462, 463

jagorii 463

platydactyla 463

Platymantis 462

plicifera 462

Hazelia 483, 484

anodon 483

picta 483

spinosa 484

Holarctic Region 441

Hoplobatrachus 469, 489, 509, 531

chinensis 469

rugulosus 469, 489, 509, 531

Hydrophylax 480, 481, 482

albotuberculata 480

everetti 481

igorata 481

luzonensis 481

tipanan 482

Hydrostentor 469

pantherinus 469

Hyla 477, 486, 487

bilineata 477

leucomystax 486

leucopogon 486

quadrilineata 486

quadrivirgata 486

sexvirgata 486

wirzi 487

Hyladactylus 474

baleatus 474

Hylaedactylus 474, 475

baleatus 474

var. *concatenata* 474

balteatus 474

bivittatus 475

celebensis 474

Holonectes (as subgenus) *conjunctus* 474

lividus 474

Hylaplesia brevipes 460

Hylarana 470, 477, 478, 479, 480, 481, 482, 489, 517, 534
 albotuberculata 480
 erythraea 477, 489, 517, 534
 everetti 481
 grandocula 478
 guttmani 478
 Hylarana (as subgenus) *varians* 482
 luzonensis 481
 mangyanum 479
 melanomenta 479
 moellendorffi 479
 nicobariensis 477
 sanguinea 482
 similis 480
 tipanan 482
Hylodes 462, 463, 472
 corrugatus 462
 Halophilus (as subgenus) 462, 463
 corrugatus 462
 dorsalis 463
 planirostris 472
Hylorana 477, 482, 486
 longipes 486
 nicobariensis 477
 varians 482

I

Ichthyophiidae 488, 489, 525, 526, 538
Ichthyophis 488, 525, 526, 538
 glandulosus 488, 525, 538
 mindanaoensis 488, 526
 monochrous 488
 weberi 488, 526
Indonesia 468, 474, 477, 487, 545
 Java 462, 463, 468, 474, 477, 487, 545
 Sumatra 545
Ingerana 461
 Ingerana mariae 461
Ingerophrynus 459, 499, 527
 philippinicus 459, 499, 527
Insecta 441
Ixalus 482, 483, 484, 485, 487
 acutirostris 484
 bimaculatus 487
 granulatus 483
 guttatus 482
 leitensis 484
 mindorensis 485
 natator 482
 var. *nubilus* 483
 nubilus 483

pictus 483
schmackeri 485

K

Kalaoula 474, 475
 conjuncta 474m 475
 conjuncta 474
 negrosensis 475
Kalophrynus 474, 514, 533
 sinensis 474, 514, 533
Kaloula 474, 475, 476, 489, 514, 516, 533, 534
 baleata 474, 514
 baleata 474
 ghoshi 474
 kalingensis 475
 conjuncta 474
 stickeli 475
 conjuncta 474, 515, 533
 meridionalis 475
 kalingensis 475, 515, 533
 kokacii 475, 515
 macrocephala 476
 negrosensis 474
 picta 475, 515, 534
 pulchra 475, 489, 516, 534
 hainana 475
 macrocephala 476
 pulchra 475
 rigida 476, 516, 534
 walteri 476, 516, 534
Kurixalus 483, 521, 536
 appendiculatus 483, 521, 536

L

Leptobrachium 473, 513, 532, 533
 lumadorum 473, 513, 532
 mangyanorum 473, 513, 533
 tagbanorum 473, 513, 533
Leptomantis 483, 487
 appendiculatus 483
 bimaculata 487
 bimaculatus 487
Limnodytes 477, 486
 celebensis 486
 erythraeus 477
Limnonectes 468, 469, 470, 471, 509, 510, 511, 512, 531, 532
 acanthi 469, 509, 531
 diuatus 469, 509
 Fejervarya (as subgenus) 468
 cancrivorus 468
 raja 468

vittiger 468
ferneri 470, 510
Hoplobatrachus (as subgenus) 468
 cancrivorus 468
 moodiei 468
 raja 468
 rugulosus 469
Limnonectes (as subgenus) 469, 470, 471
 acanthi 469
 diuatus 469
 leytensis 470, 510, 531
 macrocephalus 470
 magnus 470
 palavanensis 471
 parvus 471
 visayanus 471
 woodworthi 471
macrocephalus 470, 510, 531
magnus 470, 510, 532
micrixalus 471
palavanensis 471, 511, 532
parvus 471, 511, 532
visayanus 471, 511, 532
woodworthi 471, 512, 532
Lithobates 478, 489, 518, 535
 catesbeianus 478, 489, 518, 535

M

Macromitrium 541, 542, 543, 544, 545, 546, 547, 549
 archboldii 541, 549
 benguetense 541
 blumei 546, 549
 zollingeri 546
 eddyi **sp. nov.** 541, 542, 543, 544, 547, 549
 falcatulum 545, 549
 fasciculare 541
 formosae 541
 longicaule 545, 547, 549
 macrosporum 545, 549
 microstomum 546, 547, 549
 mindorensis 541, 549
 nepalense 541
 ochraceum 541, 545, 547, 549
 orthostichum 545, 549
 reinwardtii 546, 547, 549
 robinsonii 541
 tylostomum 541, 545, 549
 yuleanum 545, 549
 Madagascar (see under Africa) 451
 Malaysia 476, 484
 Borneo 474, 476, 484, 487
 Sarawak 476, 484, 487

Malesia 541, 542, 546, 547
Megalophrys 473
 ligayae 473
 Megophryidae 473, 489, 513, 514, 532, 533
Megophrys 473, 513, 514, 533
 ligayae 473, 513, 533
 monticola 437
 ligayae 473
 stejnegeri 473
 stejnegeri 473, 514, 533
 stejnegeri 473
 Mexico 547
Micrixalus 461, 472, 511
 diminutiva 472
 mariae 461
Microhyla 474, 476, 516, 534
 leucostigma 474
Microhyla (as subgenus) *petrigena* 476
 petrigena 476, 516, 534
 Microhylidae 474, 489, 514, 515, 516, 517, 533, 534

N

Nearctic 441, 442, 443
Nebria 441, 442, 443, 444, 445, 446, 447, 448
 baumanni **sp. nov.** 441, 442, 443, 444, 445, 446, 447, 448
 castanipes 447
 edwardsi 447
 eschschooltzii 447
 giulianii 441
 giulianii 441, 444, 445, 446, 447, 448
 lamarckensis 441, 445, 448
 meanyi 441, 444, 445, 448
 oblique 447
 ovipennis 447
 rathvoni 448
 sylvatica 441, 445, 448
 trifaria 448
 Nebriini 441
Nectophryne 460, 474
 lighti 460
 picturata 474
 New Guinea (see also Papua New Guinea) 541, 542, 545
 New Zealand 547
 North America 441, 448
Nyctixalus 483, 484, 522, 536, 537
 anodon 483
 pictus 483, 484, 522
 spinosus 484, 522, 536, 537

O

- Occidozyga* 472, 512, 532
 diminutiva 472, 512, 532
 laevis 472, 512, 532
Oceania 547
Ooeidozyga 472
 diminutives 472
 laevis 472
 laevis 472
Oreophryne 476, 517, 534
 annulata 476
 anulata 476, 517, 534
 nana 476, 517, 534
Orthotrichaceae 541, 542
Oxydozyga 472
 laevis 472
 laevis 472
Oxyglossis laevis 472
Oxyglossus laevis 472

P

- Papua New Guinea* (see also *New Guinea*) 541, 542, 545
Pelophryne 459, 460, 500, 527
 albotaeniata 459, 460, 500
 brevipes 460, 500, 527
 lighti 460, 500, 527
Philautus 463, 466, 483, 484, 485, 486, 487, 522, 523, 524, 537
 acutirostris 484, 522, 537
 anodon 483
 basilanensis 484
 bimaculatus 487
 emembranatus 486
 everetti 484, 522, 537
 hazela 463
 leitensis 484, 523, 537
 lissobrachius 485
 longicrus 485, 523, 537
 mindorensis 485
 montanus 487
 Philautus (as subgenus) 484, 486
 acutirostris 484
 emembranatus 486
 lissobrachius 485
 longicrus 485
 poecilus 485
 surdus 486
 surrufus 486
 pictus 483
 pictus 483
 poecilus 485

- poecilus* 485, 523, 537
 polillensis 466
 schmackeri 485, 523
 spinosus 484
 surdus 485, 524, 537
 surrufus 486, 524
 williamsi 485
 woodi 484
 worcesteri 486, 524, 537
 zamboangensis 487
Philippine Islands (also as *Philippines*) 457, 541
 Alabat 461, 463, 470, 472, 475
 Babuyan Island Group 475
 Balabac 459, 469, 472, 473, 480
 Basilan 460, 470, 471, 472, 473, 474, 478, 483, 484, 488
 Biliran 463, 471, 478, 483
 Bohol 460, 461, 462, 463, 467, 468, 469, 470, 471, 472, 473, 474, 475, 478, 483, 484, 485, 486, 487, 488
 Borocay 468, 475, 477
 Busuanga 459, 468, 469, 472, 480, 482, 483, 487
 Cagayan 462, 487
 Cagraray 462, 463, 468, 469, 472, 480, 487
 Calagna-an 463, 468, 471, 472, 477, 487
 Calamian Islands 469
 Calauit 472, 487
 Caluya 468, 469, 475, 487
 Camiguin 546, 549
 Camiguin Norte 472, 487
 Camiguin Sur 462, 468, 469, 470, 471, 472, 474, 475, 476, 478, 483, 487, 488
 Camotes Island Group 462
 Catanduanes 461, 462, 463, 470, 472, 475, 480, 482, 487, 488
 Cebu 461, 462, 463, 468, 469, 470, 471, 472, 475, 476, 487
 Cocomo 461, 469, 475
 Culion 459, 469, 474, 479, 480, 482, 483
 Cuyo 468, 475
 Dinagat 459, 461, 462, 463, 468, 469, 470, 471, 472, 473, 474, 475, 478, 483, 487, 488
 Dumaran 459, 468, 487
 Gigante Norte 464
 Gigante Sur 464
 Gigantes Norte 461, 468, 487
 Gigantes Sur 468, 487
 Guimaras 468, 469, 471, 472, 475, 477, 487
 Inampulugan 468, 472, 487
 Jau 468
 Lapinig Grande 468, 475
 Lapinin Chico 468

- Leyte 460, 461, 462, 463, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 480, 483, 484, 485, 487, 488
- Lubang 461, 463, 468, 469, 472, 475, 487
- Luzon 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 472, 473, 475, 476, 477, 478, 480, 481, 482, 483, 486, 487, 488, 541, 546, 547
- Albay Province 462
- Bulacan Province 461
- Cagayan Province 462
- Camarines Peninsula 464
- Isabela Province 466, 468
- Kalinga Subprovince 481
- Laguna Province 465, 472
- Mountain Province 462, 467, 475, 476
- Oriental Province 465, 473
- Quezon Province 461, 467, 476
- Tayabas Province 470
- Mactan 468, 475, 487
- Marinduque 463, 468, 469, 470, 472, 482, 487
- Masbate 461, 463, 468, 469, 470, 471, 472, 477, 487
- Mindanao 459, 460, 461, 462, 463, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 478, 480, 481, 483, 484, 485, 486, 487, 488, 541, 542, 545, 546, 547, 549
- Agusan del Norte Province 485
- Agusan Province 460, 471, 473, 478, 484
- Bukidnon Province 542, 545, 546, 547
- Davao Province 488
- Davao Oriental Province 546, 547
- Misamis Occidental Province 486
- North Cotabato Province 545, 546
- South Cotabato Province 478
- Zamboanga 459, 460, 472, 473, 481
- Zamboanga Del Sur Province 473
- Mindoro 461, 462, 468, 469, 472, 473, 475, 476, 477, 478, 479, 485, 487, 488, 546, 547
- Oriental Mindoro Province 479
- Moro 469
- Negros 461, 462, 463, 465, 467, 468, 469, 470, 471, 472, 475, 477, 487, 488
- Pacijan 468, 475, 487
- Palawan 459, 461, 468, 469, 471, 472, 473, 474, 475, 476, 477, 480, 482, 483, 484, 485, 487, 489, 546
- Pan de Azucar 463, 468, 469, 487
- Panay 461, 462, 463, 465, 466, 468, 469, 471, 472, 475, 477, 478, 487
- Antique Province 466
- Polillo 461, 462, 463, 465, 466, 468, 469, 470, 472, 475, 476, 480, 482, 487, 488
- Ponson 462, 468, 475
- Poro 462, 471, 475
- Rapu-Rapu 462, 463, 468, 470, 475, 480
- Romblon Island Group 461, 464, 468, 469, 470, 471, 472, 475, 477, 487, 488
- Samar 460, 461, 462, 463, 467, 470, 471, 472, 473, 474, 475, 476, 477, 478, 480, 483, 484, 485, 487, 488
- Eastern Samar Province 461
- Semirara 473, 475, 479, 487
- Sibay 487
- Sibuyan 467
- Sicogan 487
- Sicogon 461, 462, 463, 468, 471, 472
- Siquijor 462, 471, 475, 488
- Sulu Archipelago 468, 470, 472, 474, 475, 477, 479, 487
- Jolo 487
- Papahag Island 479
- Tawi-tawi 476
- Tablas 464
- Ticao 461, 463, 468, 471
- Tintiman 468
- Verde 461, 468, 475, 487
- Visayan Region 546, 547
- Phrynixalus* 476
- annulatus* 476
- anulatus* 476
- Phrynoglossus* 472
- diminutiva* 472
- diminutives* 472
- laevis* 472
- laevis* 472
- Phrynoidis* 460
- agua* 460
- Platymantis* 461, 462, 463, 464, 465, 466, 467, 468, 501, 502, 503, 504, 505, 506, 507, 508, 528, 529, 530, 531, 539
- banahao* 461, 501, 528
- bayani* 461, 501, 528
- biak* 461, 501, 528
- cagayanensis* 462, 528
- cagayensis* 501
- cornuta* 462
- cornutus* 462, 502, 528
- corrugata* 462
- corrugatus* 462, 502, 528
- corrugatus* 462
- diesmosi* 462, 502, 528
- dorsalis* 463, 502, 528
- guentheri* 463, 465, 503, 529
- hazela* 463, 503, 529

indepressa 464
indepressus 464, 503
ingeri 463
insulata 464
insulatus 464, 503, 529
isarog 464, 504, 529
Lahatnanguri (as subgenus) 461, 462, 464, 466
 biak 461
 cagayanensis 462
 cornutus 462
 insulatus 464
 levigatus 464
 pygmaeus 466
lawtoni 464, 504, 529
levigata 464
levigatus 464, 504, 529
Lupacolus (as subgenus) 462, 463, 464, 465, 466, 467, 468
 dorsalis 463
 indepressus 464
 mimulus 465
 naomii 465
 paengi 466
 pseudodorsalis 466
 quezoni 467
 spelaeus 467
 taylori 468
luzonensis 465
meyeri 463
mimula 465
mimulus 465, 505
Montana 465
montanus 465, 505, 529
naomiae 465
naomii 465, 505
negrosensis 465, 505, 530
paengi 466, 506, 530
panayensis 466, 506
plicifera 462
polillensis 466, 506, 530
polilloensis 466
pseudodorsalis 466, 506
pygmaea 466
pygmaeus 466, 507, 530
quezoni 467, 539
rabori 467, 507, 530
reticulatus 464
sierramadrensis 467, 507, 530
spelaea 467
spelaeus 467, 507, 530
subterrestris 467, 508, 530
Tagomukhus (as subgenus) *corrugatus* 462

Tahananpuno (as subgenus) 461, 462, 463, 465, 467
 banahao 461
 diesmosi 462
 guentheri 463
 luzonensis 465
 negrosensis 465
 rabori 467
taylori 468, 508, 531
Tirahanulap (as subgenus) 463, 464, 465, 466, 467
 hazela 463
 montanus 465
 panayensis 466
 polillensis 466
 sierramadrensis 467
 subterrestris 467
Plectropus 474, 475
 baleatus 474
 pictus 475
Polypedates 480, 483, 484, 485, 486, 487, 488, 524, 525, 538
 everetti 484
 hecticus 486
 leucomystax 486, 487, 524, 538
 linki 487
 macrotis 487, 525, 538
 maculatus quadrilineatus 486
 pardalis 488
 quadrilineatus 486
 rugosus 486
 similis 480
 surdus 485
Pulchrana 478, 479, 480, 518, 519, 535
 grandocula 478, 518
 guttmani 478
 mangyanum 479, 518, 535
 melanomenta 479, 518
 moellendorffi 479, 519, 535
 similis 480, 519, 535

R

Rana 460, 462, 463, 468, 469, 470, 471, 477, 478, 479, 480, 481, 482, 483
 Aquarana catesbeiana 478
 acanthi 469
 albotuberculata 480
 brasiliانا 460
 burkilli 469
 cancrivora 468
 cancrivora 468
 raja 468

- catesbeiana* 478
catesbyana 478
Chalcorana (as subgenus) 480, 481
 lbotuberculata 480
 everetti 481
 luzonensis 481
chinensis 469
conspersa 478
diuata 469
dubita 480
erythraea var. *elongate* 477
esculenta chinensis 469
Euphlyctis (as subgenus) 468
 cancrivora 468
 magna 470
 magna macrocephala 470
 microdisca parva 471
 rugulosa 469
everetti 480
 albotuberculata 480
 everetti 481
 luzonensis 481
Fejervarya (as subgenus) 468
 vittiger 468
 vittigera 468
gigas 460
grandocula 478
guerreroi 481
guttatus 483
humeris-armata 460
Hylarana (as subgenus) 479, 480, 481, 482
 everetti 480, 481
 luzonensis 481
 mearns 480
 moellendorffi 479
 sanguinea 482
 signata 480
 varians 482
Hylorana (as subgenus) 477, 481
 erythraea 477
 nicobariensis 477
igorata 481
javanica 477
lemniscata 477
leytensis 470
limnocharis 469
 vittigera 468
Limnodytes (as subgenus) *erythraea* 477
luzonensis 481
macrodon 470, 471
 acanthi 469
 blythii 469, 470, 471
 macrocephala 470
 magna 470
 visayanus 471
macularia var. *javanica* 477
magna 470, 471
 acanthi 469
 macrocephala 470
 magna 470
 visayanus 471
mangyanum 479
marina 460
maxima 460
mearnsi 480
melanomenta 479
merilli 481
merrilli 480
micrixalus 471
microdisca 470
 leytensis 470
 palavanensis 471
 parva 471
mindanensis 470
modesta 470
 magna 470
moellendorffi 479
moodiei 468
mugicus 478
mugiens 478
nantaiwuensis 478
nicobariensis 477
 javanica 477
 nicobariensis 477
Novirana, Aquarana catesbeiana 478
palavanensis 471
parva 471
philippinensis 478
pipiens 478
Platymantis (as subgenus) *dorsalis* 463
Pulchrana (as subgenus) 479, 480
 melanomenta 479
 moellendorffi 479
 similis 480
raja 468
Rana (as subgenus) 468
 catesbeiana 478
 crancrivora 468, 478
rugulosa 469
sanchezi 477
sanguinea 482
Sanguirana (as subgenus) 482
 sanguinea 482
 varians 482

scapularis 478
signata 478
signata 478, 479, 480
 gradocula 478
 moellendorffi 479
 similis 480
similis 480
suluensis 477
Sylvirana (as subgenus) *nicobariensis* 477
tafti 481
taurina 478
tigerina var. *burkilli* 469
tigrina 468, 469
 angustopalmata 468
 cancrivora 468
 pantherina 469
 rugulosa 469
 var. *cancrivora* 468
tipanan 482
varians 482
vittigera 468
woodworthi 471
yakani 478
Ranidae 477, 489, 517, 518, 519, 520, 534, 535, 536
Rhacophoridae 483, 489, 521, 522, 523, 524, 525, 536, 537, 538
Rhacophorus 463, 466, 483, 484, 485, 486, 487, 488, 525, 538
 anodon 483
 appendiculatus 483
 appendiculatus 483
 chasen 483
 bimaculatus 487, 525, 538
 chasen 483
 emembranatus 486
 everetti 484
 granulosus 483
 hecticus 486
 kampeni 487
 leprosus spinosus 484
 Leptomantis (as subgenus) *bimaculatus* 487
 leucomystax 486
 leucomystax 486
 linki 487
 quadrilineata 4867
 quadrilineatus 486
 var. *sexvirgata* 486
 lissobrachius 485
 macrotis 487
 maculatus 486
 himalayensis 487
 leucomystax 487
 var. *quadrilineata* 486
pardalis 488, 525, 538
 pardalis 488
 pulchellus 488
 rhysocephalus 488
Philautus (as subgenus) 463
 hazela 463
 acutirostris 484
 alticola 487
 anodon 483
 basilanensis 484
 bimaculatus 487
 leitensis 485
 mindorensis 485
 pictus 483
 schmackeri 485
 spinosus 484
 williamsi 485
 woodi 484
phyllopygus 483
polillensis 466
Polypedates (as subgenus) 487
 leucomystax 487
 macrotis 487
 quadrilineatus 487
pulchellus 488
Rhacophorus (as subgenus) 483, 484, 485, 487
 appendiculatus appendiculatus 483
 buergeri 484, 485
 everetti 484
 surdus 485
 chasen 483
 everetti 484
 hecticus 487
 himalayanus 487
 kampeni 487
 leucomystax leucomystax 487
 lincki 487
 macrotis 487
 pardalis 488
 pulchellus 488
 surdus 485
 wirzi 487
 rhysocephalus 488
 rizali 488
 surdus 485
Rhinella 460, 461, 489, 500, 527
 marina 460, 461, 489, 500, 527
 marinus 461

S

Sanguirana 480, 481, 482, 519, 520, 521, 535, 536

albotuberculata 480, 519, 535
aurantipunctata 480, 519, 535
everetti 480, 481, 520, 535
igorota 481, 520, 536
luzonensis 465, 481, 504, 520, 529, 536
sanguinea 482, 520, 536
tipanan 482, 521, 536
varians 482
Socotra 451
Solomon Islands 545
Sphenophryne 474
 beyeri 474
 fusca 474
 leucostigma 474
Staurois 472, 482, 483, 521, 536
 diminutives 472
 natator 482, 483, 521, 536
 nubilus 483, 521, 536
Sylvirana 477
 nicobariensis 477

T

Taylorana 461
 mariae 461
Tigrina 469
 rugulosa 469

U

United States 441
 Alaska 445
 California 441
 Mono County 445
 Sierra Nevada 445
 Siskiyou County 445
 Nevada 441, 442, 443, 444, 445, 447
 Clark County 441, 443, 444, 445, 448
 Esmeralda County 441
 Washington 445

CALIFORNIA ACADEMY OF SCIENCES

PROCEEDINGS SERIES

INSTRUCTIONS TO AUTHORS

Authors planning to submit papers for consideration for publication in the Academy's *Proceedings*, *Occasional Papers*, or *Memoir* series must follow the directions given below in preparing their submissions. Under some circumstances, authors may not be able to comply with all the computer-based requirements for submission. Should this be the case, please contact the Editor or Associate Editor for guidance on how best to present the materials.

The Scientific Publications Office of the Academy prepares all materials for publication using state-of-the-art, computer-assisted, page-description-language software. Final copy is sent to the printer for printing. The printer does not modify the files sent for printing. Therefore, it falls to the authors to check carefully page proof when it is returned for approval. Ordinarily, all communication with authors is done via email and galley and page proofs of manuscripts, including figures, are transmitted as attachments to email communications. Again, exceptions to this will be made in the event that an author is unable to communicate in this way.

Authors are expected to provide digital copies of both manuscript text files and images, as well as a paper printout of their manuscript. Please note the following:

TEXT: Text can be in Microsoft Word, as a Word document file, WordPerfect, also as a WP document file, or, best of all, as an "rtf" (rich text format) file, which can be produced by most word processors. Authors who use non-standard fonts must include file copies of those fonts so that their symbols can be reproduced accurately. However, it is strongly recommended that the type style "New Times Roman" be used throughout and that the Symbols and Bookshelf Symbol 1 and 3 fonts be used for such items as ♂, ♀, μ, etc. Note, words must not be typed in all capital letters either in the text or bibliography; small caps are acceptable.

IMAGES: Images should be in either JPG (JPEG), or TIF (TIFF) format. Resolution for grayscale images should be at least 600 ppi (1200 ppi if possible, especially for photomicrographs), and 300 ppi (600 ppi acceptable) for color. All images should be sized so that none exceeds a maximum print size of 5.5"×7.875" (140 mm × 200 mm).

TABLES: Our processing software allows for direct importation of tables. This reduces the chances for errors being introduced during the preparation of manuscripts for publication. However, in order to use this feature, tables must be prepared in Microsoft Excel or in Microsoft Word using Word's table feature; do not prepare tables using tabs or space bars. Complex tables not prepared as described above will be returned to the author for revision.

DIGITAL FILES: IBM or MAC formatted disks will be accepted subject to the following conditions: (a) floppy disks must not exceed 1.4 mb and (b) zip disks, preferably IBM format, must not exceed 100mb. Authors are encouraged to submit their digital files on CD-ROM (CD-R formatted disks NOT CD-RW) inasmuch as these can be read by nearly all CD-ROM drives.

FILE NAMING PROTOCOLS: To facilitate the handling of digital files submitted by authors, the following file-naming conventions are to be followed: text files should bear the author's last name (in the case of multiple authors, only the first author's name) followed by a space and a date in the format mmyy (e.g., 0603 for June 2003) to yield a file name such as **Gosliner 0603.doc** or **Williams 0603.rtf**. If an author has submitted two or more manuscripts and must distinguish between them, then the naming should include an additional numeral: **Gosliner1 0603.doc** for the first manuscript, **Gosliner2 0603.doc** (or .rtf) for the second. Figures should follow similar conventions, as follows: **Gosliner F1 0603.tif**, **Gosliner F2 0603.tif**, for figures in the first manuscript and, if more than one manuscript, then **Gosliner1 F1 0603.tif** etc. for the figures associated with the first manuscript and **Gosliner2 F1 0603.tif** etc. for those with the second. Following these conventions will insure that figures submitted by one author are always maintained distinct from those submitted by another. Tables submitted as Excel files should follow the same naming conventions except the file type designation will be ".xls": e.g., **Gosliner T1 0603.xls**. Please note that extraneous periods are omitted in file names.

BIBLIOGRAPHY FORMAT: Three bibliographic styles are accommodated in the Academy's scientific publications, one commonly used in scientific journals publishing papers in systematic and evolutionary biology, a second used mainly in the geological literature, and lastly, the format most commonly used in the humanities by historians of science. On request, the author will be sent a style sheet that includes samples of the three formats. Authors are also encouraged to examine a copy of the latest published *Proceedings*. In all instances, however, authors should not abbreviate journal names but spell them out completely. For books, the reference must include the publisher and city of publication. It is recommended that the total number of pages in the book also be given.

SUBSCRIPTIONS/EXCHANGES

The *Proceedings* series of the California Academy of Sciences is available by exchange or subscription. For information on exchanges, please contact the Academy Librarian via regular mail addressed to the Librarian, California Academy of Sciences, 55 Music Concourse Drive, Golden Gate Park, San Francisco, CA 94118 U.S.A. or via email addressed to **hyaeger@calacademy.org**. Subscription requests, including information on rates, should be addressed to Scientific Publications, California Academy of Sciences, 55 Music Concourse Drive, Golden Gate Park, San Francisco, CA 94118 U.S.A. or via email to the Editors at **aleviton@calacademy.org** or **gwilliams@calacademy.org**

Subscription price for 2010: \$75 (US) includes mailing to U.S. and Canadian addresses and \$85 to all others.

The *Occasional Papers* and *Memoirs* are not available by subscription. Each volume is priced separately. *Occasional Papers*, *Memoirs*, and individual issues of the *Proceedings* are available for purchase through the Academy's Office of Scientific Publications. Visit us on the web at <<http://research.calacademy.org/research/scipubs/>>.

COMMENTS

Address editorial correspondence or requests for pricing information to the Editor, Scientific Publications Office, California Academy of Sciences, 55 Music Concourse Drive, Golden Gate Park, San Francisco, CA 94118 U.S.A. or via email to the Editor, Scientific Publications, at **aleviton@calacademy.org** or **gwilliams@calacademy.org**

Table of Contents

DAVID H. KAVANAUGH: A New Species of *Nebria* Latreille (Insecta: Coleoptera: Carabidae: Nebrini) from the Spring Mountains of Southern Nevada441

THOMAS F. DANIEL: *Anisotes tablensis* (Acanthaceae), a New Species from Southwestern Madagascar451

ARVIN C. DIESMOS, JESSA L. WATTERS, NICHOLAS A. HURON, DREW R. DAVIS, ANGEL C. AL-
CALA, RONALD I. CROMBIE, LETICIA E. AFUANG, GENEVIEVE GEE-DAS, ROGELIO V. SISON,
MARITES B. SANGUILA, MICHELLE L. PENROD, MARIE J. LABONTE, CONNER S. DAVEY, E.
AUSTIN LEONE, MAE L. DIESMOS, EMERSON Y. SY, LUKE J. WELTON, RAFAEL M. BROWN,
AND CAMERON D. SILER: Amphibians of the Philippines, Part I: Checklist of the Species
.....457

BENITO C. TAN AND JAMES R. SHEVOCK: Species of *Macromitrium* (Orthotrichaceae) New
to the Mindanao Region and the Philippines with One Species New to Science . .541

Index551